arXiv:1011.0118v3 [math.GR] 9 Dec 2010

Space functions of groups.

A.Yu.Olshanskii *

November 8, 2011

Abstract

We consider space functions $s(n)$ of finitely presented groups $G = \langle A \mid R\rangle$. (These functions have a natural geometric analog.) To define $s(n)$ we start with a word w over A of length at most n equal to 1 in G and use relations from R for elementary transformations to obtain the empty word; $s(n)$ bounds from above the tape space (or computer memory) one needs to transform any word of length at most n vanishing in G to the empty word. One of the main results obtained is the following criterion: A finitely generated group H has decidable word problem of polynomial space complexity if and only if H is a subgroup of a finitely presented group G with a polynomial space function.

Key words: generators and relations in groups, space complexity, algorithmic word problem, van Kampen diagram

AMS Mathematical Subject Classification: 20F05, 20F06, 20F65, 20F69, 03D15, 03D40, 03D10

1 Introduction

Time and space complexities are the main properties of algorithms. Their counterparts in Group Theory are the Dehn and filling length (or space) functions of finitely presented groups. In this paper, we study the interrelation of space functions of groups and the space complexity of the algorithmic word problem in groups.

Let $G = \langle A \mid R\rangle$ be a group presentation, where A is a set of generators and R is a set of defining relators. Recall that relators belong to the free group with basis A, and a group word w in generators A (i.e., a word over $A^{\pm 1}$) represents the identity of G iff there is a *derivation*

$$w \equiv w_0 \to w_1 \to \cdots \to w_{t-1} \to w_t \equiv 1 \tag{1.1}$$

where 1 is the empty word, the sign $\equiv$ is used for letter-by-letter equality of words, and for every $i = 1, \dots, t$, the word w_i results from w_{i-1} after application of one of the elementary R-transformations. As such transformations one can take free reductions of subwords $aa^{-1} \to 1$ ($a \in A^{\pm 1}$), removing subwords $r^{\pm 1}$, where $r \in R$, and the inverse transformations.

The minimal non-decreasing function $f(n)\colon \mathbb{N} \to \mathbb{N}$ such that for every word w vanishing in G and having length $||w|| \le n$, there exists a derivation (1.1) with $t \le f(n)$, is called the Dehn function of the presentation $G = \langle A \mid R\rangle$ [13]. For *finitely presented* groups (i.e., both sets A and R are finite) Dehn functions are usually taken up to equivalence to get rid of the dependence on a finite presentation for G (see [18]). To introduce this equivalence $\sim$, we write $f \preceq g$ if there is a positive integer c such that

$$f(n) \le cg(cn) + cn \quad for \ \ any \ \ n \in \mathbb{N} \tag{1.2}$$

For example, we say that a function f is *polynomial* if $f \preceq g$ for a polynomial g. From now on we use the following equivalence for non-decreasing functions f and g on $\mathbb{N}$.

$$f \sim g \quad if \ \ both \ \ f \preceq g \ \ and \ \ g \preceq f \tag{1.3}$$

*The author was supported in part by the NSF grant DMS 0700811 and by the Russian Fund for Basic Research grant 08-01-00573

It is not difficult to see that the Dehn function $f(n)$ of a finitely presented group G is recursive (or bounded from above by a recursive function) iff the word problem is algorithmically decidable for G (see [11], [6]). In this case, the word problem can be solved by a primitive algorithm that, given a word w of length n, just checks if there exists a derivation (1.1) of length $\leq f(n)$. Therefore the nondeterministic *time* complexity of the word problem in G is bounded from above by $f(n)$. Moreover if H is a finitely generated subgroup of G, then one can use the rewriting procedure (1.1) for H, and so the nondeterministic time complexity of the word problem for H is also bounded by $f(n)$.

It turns out that a converse statement is also true. Assume that the word problem can be solved in a *finitely generated* group H by a nondeterministic Turing machine (NTM) with time complexity $T(n)$. Then H is a subgroup of a *finitely presented* group G with Dehn function equivalent to $n^2T(n^2)^4$ [4]. As the main corollary, one concludes that the word problem of a finitely generated group H has time complexity of class NP (i.e., there *exists* a non-deterministic algorithm of polynomial time complexity, that solves the word problem for H) iff H is a subgroup of a finitely presented group with polynomial Dehn function.

We want to obtain similar statements for space functions. It is clear that to perform the rewriting (1.1) one needs space equal to $\max_{0\leq i\leq t}||w_i||$, and this observation leads to the definition of space function of a finitely presented group. However when handling groups, one can enlarge the set of elementary R-transformations and obtain different definitions of space functions. One can either (1) consider only the transformations we defined above and obtain the filling length function introduced by Gromov [13] (also see [12], [3]); or (2) also allow replacement of words by their cyclic permutations as was suggested by Bridson and Riley [7]; or (3) additionaly use the replacement of a word $w \equiv uv$ by the pair (u, v) if both u and v are trivial in G (see [7] again). Starting with these different sets of transformations one comes to different space functions called in [7], respectively, filling length function (FL), free filling length functions (FFL), and fragmenting free filling length functions ($FFFL$). Each of these functions has a visual geometric interpretation in terms of the transformations of loops in the Cayley complex of G using, respectively, null-homotopy, free null-homotopy, and free null-homotopy with bifurcations. It is proved in [7] that these functions behave differently for the same finitely presented group G, for instance, $FFFL$ can grow linearly while FL and FFL have exponential growth. There are many other features of these functions presented in [7] to justify their "inclusion in the pantheon of filling invariants".

In this paper, we choose the third version ($FFFL$), and this choice is justified by the theorems on the connections between such functions and the space complexity of the word problem for groups.[1] Thus we operate with finite sequences of words $W = (w_1, \ldots, w_s)$ over a group alphabet A. Given a finitely presented group G, we say that a finite sequence $W' = (w'_1, \ldots, w'_{s'})$ results from W after application of an elementary R-transformation if $s' \in \{s-1, s, s+1\}$ and one of the following is done for some w_i $(i = 1, \ldots, s)$:

- a subword aa^{-1} is removed from or inserted to w_i $(a \in A^{\pm 1})$;
- a subword r or r^{-1} is removed from or inserted to w_i $(r \in R)$;
- w_i is replaced by a cyclic conjugate;
- $w_i \equiv uv$, and w_i is replaced by the pair u, v, i.e. $W' = (w_1, \ldots, w_{i-1}, u, v, w_{i+1}, \ldots, w_s)$;
- w_i is removed if it is empty, i.e. $W' = (w_1, \ldots, w_{i-1}, w_{i+1}, \ldots, w_s)$.

Clearly, we have $w = 1$ in the group G iff there exists an R-rewriting starting with (w) and ending with the empty string ( ).

For every finite sequence $W = (w_1, \ldots, w_s)$ we set $||W|| = \sum_{i=1}^{s} ||w_i||$. By definition, the space of a rewriting $W_0 \to \cdots \to W_t$ is $\max_{j=0}^{t} ||W_j||$. If a word vanishes in G, then $space(w) = space_G(w)$ is the minimum of spaces of all rewritings starting with (w) and ending with the empty string. The *space function* of the group presentation $G = \langle A \mid R \rangle$ (or briefly, of the group G) is the function

$$S_G(n) = \max(space(w), \ \ where \ \ w = 1 \ \ in \ \ G \ \ and \ \ ||w|| \leq n)$$

[1] An embedding statement for the FL-functions was conjectured by J.-C. Birget in [3].

The space functions of finitely presented groups will be regarded up to the equivalence defined by (1.2) and (1.3), and so their growth will be at least linear. It is observed in [7] that the equivalence class of S_G does not depend on a finite presentation of the group G, moreover this class is invariant under quasi-isometries.

An accurate definition of the space complexity (function) $f(n)$ for a Turing machine (TM) will be recalled in Subsection 2.1. Now we just note that it is conventional that for a multi-tape TM, the function $f(n)$ counts only the space of work tapes. However since the space complexities of machines are taken here up to the same equivalence as the space functions of groups, the adding of the space of the input tape does not change the equivalence class.

The sequence $W_0 \to \cdots \to W_t$ can be easily produced by an NTM such that the computation needs at most $2\max_{i=1}^{t} ||W_i|| + const$ tape squares. (See also Section 3 of [27] or Remark 2.4 in [7].) This immediately implies

Proposition 1.1. *The space function of a finitely presented group G is equivalent to the space complexity of a non-deterministic two-tape TM. The language accepted by this machine coincides with the set of words equal to* 1 *in the group.*

In particular, the non-deterministic space complexity of the word problem in a finitely presented group G does not exceed the space function of G. It follows from [18], [8] that the literal converse statement fails. Moreover, a counter-example can be given by Baumslag's 1-relator group [1] $G = \langle a, b \mid (aba^{-1})b(aba^{-1})^{-1} = b^2 \rangle$ because the space function of G is not bounded from above by any multi-exponential function (see papers of S.Gersten [10] and A.Platonov [25]) while the space complexity of the word problem for G is at most exponential. This was proved by M. Kapovich and Schupp (unpublished), moreover, it is polynomial (announced by A. G. Myasnikov, A. Ushakov, and Dong Wook Won). Therefore the correct formulation has to take into consideration that the algorithm from Proposition 1.1 solves the word problem not only for G but also for every finitely generated subgroup of the group G. In the deterministic case we get a sharper formulation:

Theorem 1.2. *Let H be a finitely generated group such that the word problem for H is decidable by a deterministic TM (DTM) with space complexity $f(n)$. Then H is a subgroup of a finitely presented group G with space function equivalent to $f(n)$.*

One can choose the group G so that $f(n)$ is also equivalent to the logarithm of the Dehn function of G.

The main corollary of this theorem applies to polynomial space complexity. We say that a finitely generated group G belongs to the class $PSPACE$ (to $NPSPACE$) if the word problem for G is decidable by some DTM (some NTM) with polynomial space complexity. But $NPSPACE = PSPACE$ by the remarkable theorem of Savitch (see [9], Corollary 1.31) (Since there is no similar equality for deterministic *time* complexity, it therefore has no natural algebraic counterpart.) Therefore in contrast to the main result of [4], Proposition 1.1 and Theorem 1.2 eliminate any non-determinism in

Corollary 1.3. *A* finitely generated *group H belongs to $PSPACE$ iff H is a subgroup of a* finitely presented *group G having polynomial space function.*

Thus, given a 'good' algorithm solving the word problem in H (e.g., using a matrix representation of H, etc.), it is possible to find a bigger group G whose deterministically modified ('silly') natural algorithm solves the word problem for both H and G, and whose space complexity is not much worse than the space complexity of the original algorithm. Note that most common finitely generated groups belong to $PSPACE$, with a wide range from linear groups (two-tape machines need just logspace on the work tape to solve their word problem [16]) to free Burnside groups of large odd exponents (N.Boatman, unpublished).

Another natural question raised in this paper is the realization problem: Which functions $f(n)\colon \mathbb{N} \to \mathbb{N}$ are, up to equivalence, the space functions of finitely presented groups? There are not many examples; linear and exponential ones can be found in [7], but it is not easy even to specify a group with space function n^2.

Theorem 1.4. *The space complexity $f(n)$ of arbitrary DTM M is equivalent to the space function of some finitely presented group G.*

In addition, one can choose the group G so that $f(n)$ is also equivalent to the logarithm of the Dehn function of G, and there is a one-to-one linear time reduction of the decision problem of M to the word problem of G.

This theorem gives a tremendous class of space functions of groups, including functions equivalent to $[\exp\sqrt{n}]$, $[n^k]$ ($k \in \mathbb{N}$), $[n^k \log^l n]$, $[n^k \log^l(\log\log n)^m]$, etc. Note that we do not assume in the formulation of Theorem 1.4 that the function $f(n)$ is superadditive (i.e., $f(m+n) \geq f(m)+f(n)$) or grows sufficiently fast. (Compare with Theorem 1.2 [27] on Dehn functions of groups.) It follows, in particular, that there exists a finitely presented group whose space function is not equivalent to any superadditive function. Recall that it is unknown if the Dehn function of arbitrary finitely presented group is equivalent to a superadditive function; see [15].

J.-C. Birget observed (email communication) that in addition, Theorem 1.4 implies a statement that does not involve space functions of groups (see also Remark 5.9.) The first claim of Corollary 1.5 below follows, in fact, from each of the papers [28] (by B.A.Trakhtenbrot) and [29] (by M.K. Valiev). Let us start with a DTM M solving a $PSPACE$ complete problem (for the definition and the existence see [9], Theorem 3.23). Then choose a finitely presented group $G = G(M)$ in accordance with Theorem 1.4, i.e., G has a polynomial space function. By Proposition 1.1, G belongs to $NPSPACE = PSPACE$, and so we have

Corollary 1.5. *There is a finitely presented group G with $PSPACE$ complete word problem. Moreover the group G can be chosen with polynomial space function.*

Theorem 1.2 implies a non-deterministic corollary. To formulate it we recall that a function $f : \mathbb{N} \to \mathbb{N}$ is called *fully space-constructible* (FSC) if there exists a two-tape DTM that halts on any input x of length n after visiting exactly $f(n)$ tape squares of the work tape. Most common functions are FSC (see [9]).

Corollary 1.6. *Let H be a finitely generated group such that the word problem for H is decidable by an NTM having FSC space complexity $f(n)$. Then H is a subgroup of a finitely presented group G with space function equivalent to $f(n)^2$.*

Finally, we describe the functions n^α which are (up to equivalence) space functions of groups. Our approach is to modify the proof of Savitch's theorem from [9] and the proof from [27], where the similar problem was considered for Dehn functions if $\alpha \geq 4$, and close necessary and sufficient conditions were obtained. (See also a dense series of examples with $\alpha \geq 2$ presented by Brady and Bridson [5].) Now we have $\alpha \geq 1$ in Corollary 1.7 below. Also it is remarkable that for space functions the necessary and sufficient conditions just coincide. To formulate the criterion, we call a real number α *computable with space* $\leq f(m)$ if there exists a DTM which, given a natural number m, computes a binary rational approximation of α with an error $O(2^{-m})$, and the space of this computation $\leq f(m)$.

We have obtained the following criterion.

Corollary 1.7. *For a real number $\alpha \geq 1$, the function $[n^\alpha]$ is equivalent to the space function of a finitely presented group iff α is computable with space $\leq 2^{2^m}$.*

It follows that functions $[n^\pi]$, $[n^{\sqrt{e}}]$, and $[n^\alpha]$ with any algebraic $\alpha \geq 1$ are all space functions of finitely presented groups.

The space function is defined for a simply connected geodesic metric space under some weak restrictions, in particular, for the universal cover of any closed connected Riemannian manifold. (See [7] for details; we just note here that to define the space ($= FFFL$) function, one should consider free homotopy with the possibility of separating a loop into two loops in a bifurcation point.) It is proved in [7] (Theorem E) that if a finitely presented group G acts properly and cocompactly by isometries on such a space X, then the space function of G is equivalent to the space function of X. Since every finitely presented group is a fundamental group of a connected compact Riemannian manifold, we can use corollaries 1.4 and 1.7 to formulate one more

Corollary 1.8. *For every space complexity $f(n)$ of a DTM, there exists a closed connected Riemannian manifold M such that the space function of the universal cover $\tilde{M}$ is equivalent to $f(n)$.*

In particular, if a real number $\alpha \geq 1$ is computable with space $\leq 2^{2^m}$, then there exists such a universal cover $\tilde{M}$ with space function equivalent to n^α.

To some extent, our constructions can be traced back to the works of P.Novikov, Boon, Britton and other authors who invented group-theoretical interpretations of TM-s (see [26], ch. 12). The hub relator is a multiple copy of the accept configuration of a machine. Then we use the language of van Kampen diagrams and construct a disc diagram for arbitrary accepting computation. A computational disc has the hub cell in the center, and this hub is surrounded by a number of similar sectors. At first we have to estimate the sizes of computational discs, and to do this we should know the *generalized* space complexity of a machine. It estimates the space of computations starting with *arbitrary* accept configuration, not only with input ones. So we should modify the initial machine to be able to control the generalized space complexity. (The modification from [27] helps to control the time complexity but corrupts the space complexity.)

The next modification is due to the symmetry of algebraic relations: since $u = v$ always implies $v = u$, the algebraic version of a machine M always interprets the symmetrization of M. Thus we are concerned that the symmetrization preserves the basic characteristics, e.g. the accepted language and the space functions. We are able to do this only if the initial machine is deterministic or can be transformed to a deterministic machine under the control of basic properties. (The known symmetrization trick from [2] or [27] does not work here since it does not preserve the space complexity.) This causes the restrictions in the formulations of Theorem 1.2 and Corollary 1.6.

The interpretation problem for groups remains much harder than for semigroups even after modifying the machine because the group theoretic simulation can execute unforeseen computations with non-positive words. Boon and Novikov secured the positiveness of admissible configurations in discs with the help of an additional 'quadratic letter' (see [26], ch.12), but this involves a difficult control of parameters for the constructed group. A new approach was suggested in [27]. Invented by Sapir, S-machines can work with non-positive words on the tapes. Here we also construct an S-machine which is a somewhat modified composition of a convenient Turing machine with an 'adding machine' $Z(A)$ introduced in [24]. Fortunately, $Z(A)$ does not change the space of computations but controls positiveness of configurations.

Recall that we should not just simulate the work of a machine but construct an embedding of given group H into a finitely presented group G in the spirit of the Higman Embedding Theorem. (See [26], ch.13). To this end we use a version of the two-disc scheme applied in [23]. One can say more simply that the configurations on the boundary of discs of the first type are longer than the words written on the boundary of corresponding discs of the second type, and the surpluses are relations of the group H. Since every relation of H holds in G, we obtain a homomorphism $H \to G$ that turns out to be injective.

To estimate the space function of the group G we use van Kampen diagrams and induct on the number of hubs. The basic problem is to cut a diagram Δ having at least two hubs into two subdiagrams with hubs, so that the perimeters of the subdiagrams do not exceed the perimeter of Δ. We have to introduce a new metric where the length of a word depends on its syllable factorization. To find a shortcut we use the mirror symmetry of sectors in discs, but unfortunately for the two-disc scheme, one of the sectors has no (mirror) copies, which creates technical obstacles.[2] We study both exact and non-accurate copying for various types of complete and incomplete sectors. A number of concepts, e.g. bands, trapezia, discs were introduced in algorithmic group theory early on, in papers [27], [20], [4] and subsequent ones, and we reproduce them in Section 3 (and partly in Sections 2 and 4) in a modern form. Some others (in particular, replica, unfinished diagram) are new.

In future work, we mean to consider similar problems for semigroups where the simulating of machines is easier, and where the definition of space function does not need cyclic shifts and fragmentation of words, i.e., the space functions are FL-functions. For groups, an FL-analogue of the results from this paper is still unachievable.

[2]Note that one can give shorter proofs of Theorem 1.4 and Corollary 1.7 which do not need the two-disc scheme; but here we just obtain these results after Theorem 1.2 is proved.

2 Machines

Suppose a DTM M of space complexity $S(n)$ solves the word problem in a finitely generated group H. Then one can introduce a finitely presented group G whose relations simulate the work of M and construct a natural homomorphism $H \to G$. This will be done in Section 3. But as mentioned in the Introduction, one can prove neither that this homomorphism is injective nor that the space function of G is bounded by $S(n)$, unless M enjoys some additional properties. Therefore the machine M has to be amended at the beginning. In this section we substitute M for a non-deterministic S-machine $\mathcal{S}$ which is needed for the injectivity of the mapping $H \to G$, and prove that $\mathcal{S}$ accepts the same language and has the same space complexity as the original M. Moreover, the main Lemma 2.10 of the section claims that the generalized space complexity of $\mathcal{S}$ does not exceed its space complexity; this property will be exploited in Subsection 5.2 in order to get an upper estimate of the space function of G.

2.1 Definitions

We will use a model of recognizing TM-s which is close to the model from [27]. Recall that a *(multi-tape) TM with k tapes and k heads* is a tuple

$$M = \langle X, Y, Q, \Theta, \vec{s}_1, \vec{s}_0 \rangle$$

where X is the input alphabet, $Y = \sqcup_{i=1}^{k} Y_i$ is the tape alphabet, $Y_1 \supset X$, $Q = \sqcup_{i=1}^{k} Q_i$ is the set of states of the heads of the machine, Θ is a set of transitions (commands), $\vec{s}_1$ is the k-vector of start states, $\vec{s}_0$ is the k-vector of accept states. ($\sqcup$ denotes the disjoint union.) The sets X, Y, Q, Θ are finite.

We assume that the machine normally starts working with states of the heads forming the vector $\vec{s}_1$, with the head placed at the right end of each tape, and accepts if it reaches the state vector $\vec{s}_0$. In general, the machine can be turned on in any configuration and turned off at any time.

A *configuration* of tape number i of a TM is a word uqv where $q \in Q_i$ is the current state of the head, u is the word to the left of the head, and v is the word to the right of the head. A tape is *empty* if u, v are empty words.

A *configuration* U of the machine M is a word

$$\alpha_1 U_1 \omega_1 \alpha_2 U_2 \omega_2 ... \alpha_k U_k \omega_k$$

where U_i is the configuration of tape i, and α_i, ω_i are special separating symbols. For unification of notation, we shall treat α_i, ω_i as heads of the machine too. These heads correspond to tapes that are always empty and do not change during a computation.

An *input configuration* is a configuration where all tapes except the first one are empty, the configuration of the first tape (let us call it the *input tape*) is of the form uq, $q \in Q_1$, u is a word in the alphabet X, and the states form the start vector $\vec{s}_1$. The *accept configuration* is the configuration where the state vector is $\vec{s}_0$, the accept vector of the machine, and all tapes are empty. (The requirement that the tapes must be empty is often removed for auxiliary machines which are used in construction of bigger machines.)

To every $\theta \in \Theta$, there corresponds a command (marked by the same letter θ), i.e., a pair of sequences of words $[V_1, ..., V_k]$ and $[V_1', ..., V_k']$ such that for each $j \leq k$, either both V_j and V_j' are configurations of the tape number j, or $V_j \equiv \alpha_j q$ and $V_j' \equiv \alpha_j q'$, or $V_j \equiv q\omega_j$ and $V_j' \equiv q'\omega_j$, or $V_j \equiv \alpha_j q \omega_j$ and $V_j' \equiv \alpha_j q' \omega_j$ $(q, q' \in Q_j)$.

In order to execute this command, the machine checks if V_i is a subword of the configuration of tape i for each $i \leq k$, and if this condition holds the machine replaces V_i by V_i' for all $i = 1, \ldots, k$. Therefore we also use the notation: $\theta : [V_1 \to V_1', \ldots, V_k \to V_k']$.

Suppose we have a sequence of configurations $w_0, ..., w_t$ and a word $h \equiv \theta_1 \ldots \theta_t$ in the alphabet Θ, such that for every $i = 1, ..., t$ the machine passes from w_{i-1} to w_i by applying the command θ_i. Then the sequence $(w_0 \to w_1 \to \cdots \to w_t)$ is said to be a *computation with history h*. In this case we shall write $w_0 \cdot h = w_t$. The number t will be called the *time* or *length* of the computation.

A configuration w is called *accepted* by a machine M if there exists at least one computation which starts with w and ends with the accept configuration. We do not only consider deterministic TMs, for

example, we allow several transitions with the same left side. Moreover, for non-deterministic TMs, one may correspond identically equal executions to different symbols $\theta, \theta' \in \Theta$.

A word u in the input alphabet X is said to be *accepted* by the machine if the corresponding input configuration is accepted. The set of all accepted words over the alphabet X is called the *language* $\mathcal{L}_M$ *recognized by the machine* M.

Let $|w_i|_a$ $(i = 0, ..., t)$ be the number of tape letters (or tape squares) in the configuration w_i. (As in [27], the tape letters are called a-letters.) Then the maximum of all $|w_i|_a$ will be called the *space of computation* $C : w_0 \to w_1 \to \cdots \to w_t$ and will be denoted by $space_M(C)$. By $space_M(w)$, we denote the minimal natural number s such that there is an accepted computation of space at most s, starting with the configuration w. If $u \in \mathcal{L}_M$, then, by definition, $space_M(u)$ is the space of the corresponding input configuration w.

The number $S(n) = S_M(n)$ is the minimum of the numbers $space(u)$ over all words $u \in \mathcal{L}_M$, with $||u|| \leq n$. The function $S(n)$ will be called the *space complexity* of the Turing machine.

The definition of the *generalized space complexity* $S'(n) = S'_M(n)$ is similar to the definition of space complexity but we consider arbitrary accepted configurations w with $|w|_a = n$, not just input configurations as in the definition of $S(n)$. It is clear that $S(n) \leq S'(n)$.

To obtain the definitions of $time_M(w)$, $time_M(u)$, *time complexity* $T_M(n)$ and *generalized time complexity* $T'_M(n)$, one should replace 'space' by 'time' in the previous definitions.

One may assume that only input configurations involve the state letters from $\vec{s}_1$ and only one command is applicable to the input configurations. (Indeed, given an NTM M, one can add additional states and add new commands changing the states from the new vector $\vec{s}_1$.) Similarly, one may assume that there is a unique accept configuration $\vec{s}_0$ and a unique accepting command. Under both of these assumptions, we will say that the machine satisfies the $\vec{s}_{10}$-*condition*. These assumptions change neither the language $\mathcal{L}$ nor the functions $S_M(n)$ and $S'_M(n)$.

2.2 Machines with equivalent space and generalized space complexities

In this subsection, we construct an NTM M_2 which depends on an NTM M_1, and prove Lemma 2.1 that allows us to replace the original machine M_1 by a machine M_2 inheritting the basic characteristics of M_1 and having equivalent generalized space complexity and space complexity.

Let an NTM M_1 have k tapes, and let its first tape be the input tape. Then we add a tape numbered $k + 1$, which is empty for input configurations, and we organize the work of the 3-stage machine M_2 as sequential work of the following machines M_{21}, M_{22}, and M_{23}.

The machine M_{21} uses only one command θ_* that does not change states and adds one square with an auxiliary letter $*$ to the $(k + 1)$-st tape, i.e the command θ_* has the form

$$[q_1 \to q_1, \ldots, q_k \to q_k, q_{k+1} \to * \, q_{k+1}]$$

The machine M_{21} can execute this command arbitrarily many times while the tapes numbered $1, \ldots, k$ keep the copy of an input configuration of M_1 unchanged. Then a connecting rule $\theta_{12} : [q_1 \to q'_1, \ldots, q_k \to q'_k, q_{k+1} \to q'_{k+1}]$ changes all states of the heads and switches on the machine M_{22}.

The work of M_{22} on the tapes with numbers $1, \ldots, k$ copies the work of M_1. But the extension θ' of every command θ of M_1 to the $(k + 1)$-st tape is defined so that its application does not change the current space. More precisely, if the application of θ inserts m_1 tape squares and deletes m_2 tape squares on k tapes, then θ' inserts $m_2 - m_1$ (deletes $m_1 - m_2$) squares with letter $*$ on the $(k + 1)$-st tape if $m_1 - m_2 \leq 0$ (if $m_1 - m_2 \geq 0$). That is the $(k + 1)$-st component of θ' has the form $q'_{k+1} \to *^{m_2 - m_1} q'_{k+1}$ (resp., $*^{m_1 - m_2} q'_{k+1} \to q'_{k+1}$). Note that one cannot apply θ' if $m_1 - m_2$ exceeds the current number of squares on the tape numbered $k + 1$.

The connecting command θ_{23} is applicable when M_{22} obtains the accept configuration on the first k tapes. It changes the states and switches on the machine M_{23} erasing all squares on the $(k + 1)$-st tape (one by one), and then M_2 accepts. (The tape alphabet of M_{23} has only one letter $*$.)

Let w be a configuration of the machine M_2 such that $w \cdot \theta_*$ is defined, or such that w is obtained after an application of the connecting command θ_{12}. Then we have an input configuration on the tapes with numbers $1, \ldots, k$ (plus several $*$-s on the $(k + 1)$-st tape). We will denote by $u(w)$ the input word

u written on the first tape. It is an input word for the machine M_1 as well, and if it is accepted by M_1, the expression $space_{M_1}u(w)$ makes sense.

The connecting commands θ_{12} and θ_{23} are not invertible in M_2 by definition. Therefore every non-empty accepting computation of M_2 has history of the form $h_1h_2h_3$, or h_2h_3, or h_3, where h_l is the history for M_{2l}, $(l = 1, 2, 3)$. (To simplify notation we attribute the command θ_{12} (the command θ_{23}) to h_2 (to h_3).)

Lemma 2.1. *(a) The machines M_1 and M_2 recognize the same language $\mathcal{L}$. (b) The space complexity $S_{M_2}(n)$ and the generalized space complexity $S'_{M_2}(n)$ of M_2 are both equivalent to $S_{M_1}(n)$. (c) If w is an accepted configuration for M_2, and the command θ_* is applicable to w, then $space_{M_2}(w) = \max(space_{M_1}(u(w)), |w|_a)$.*

Proof. Assume that $u \in \mathcal{L} = \mathcal{L}_{M_1}$. Then $u \in \mathcal{L}' = \mathcal{L}_{M_2}$ because the machine M_{21} can insert sufficiently many squares (equal to $space_{M_1}(u) - ||u||$) so that the accepting computation of M_1 can be simulated by M_{22}. Also it is clear from the definition of M_2, that every accepting computation for M_2 having a history $h_1h_2h_3$ as above, simulates, at stage 2, an accepting computation of M_1 with history h_2. Therefore $\mathcal{L}' = \mathcal{L}$ and $S_{M_1}(n) = S_{M_2}(n)$.

Assume now that $C : w = w_0 \to \cdots \to w_n$ is an accepting computation of M_2 with $space_{M_2}(C) = space_{M_2}(w)$ and $h \equiv h_1h_2h_3$ is the history with the above factorization (h_1 or h_1h_2 can be empty here). If the word h_1 is empty, then $||w_0|| \geq \cdots \geq ||w_n||$ by the definition of the machines M_{22} and M_{32}. Hence the space of this computation is equal to $|w|_a$ [3]. Then let h_1 be non-empty. It follows that the machine M_2 starts working with a copy of an input configuration of the machine M_1, i.e., the input tape of this configuration contains an input word $u = u(w)$, and the additional $(k+1)$-st tape has m squares for some $m \geq 0$. Moreover $u \in \mathcal{L}$ since the computation of M_2 is accepting. We consider two cases.

Case 1. Suppose $m \geq space_{M_1}(u) - ||u||$. This inequality says that the additional tape has enough squares to enable M_{22} to simulate the accepting computation of M_1 with the input word u. Hence there is an M_2-computation $w_0 \to \cdots \to w_{n'}$ with history of the form $h'_2h'_3$, and so its space, as well as the space of our original accepting computation, is $|w|_a$.

Case 2. Suppose $m < space_{M_1}(u) - ||u||$. Then there is a computation $w_0 \to \cdots \to w_{n'}$ such that the commands of its M_{21}-stage insert squares until the total number of squares of the $(k+1)$-st tape becomes equal to $space_{M_1}(u) - ||u||$, and then the machines M_{21} and M_{23} work in their standard manner. The space of this (and the original) computation is $space_{M_1}(u)$.

The estimates obtained in cases 1 and 2 prove statement (c) of the lemma. They also show that

$$S_{M_1}(n) = S_{M_2}(n) \leq S'_{M_2}(n) \leq \max(S_{M_1}(n), n)) \sim S_{M_1}(n)$$

and statement (b) is completely proved too. □

2.3 Symmetric machines

A simulation of the work of a machine M by algebraic relations leads, in fact, to the simulation of the machine M^{sym} capable of inverting every command of M. Therefore we must control the basic properties of such symmetrization, and this will be done in Lemma 2.4.

For every command θ of a TM, given by the vector $[V_1 \to V'_1, \ldots, V_k \to V'_k]$, the vector $[V'_1 \to V_1, \ldots, V'_k \to V_k]$ also gives a command of some TM. These two commands θ and θ^{-1} are called *mutually inverse*.

From now on we will assume that the machine M_1 we started with in Subsection 2.2 is a DTM and satisfies the $\vec{s}_{10}$-condition.

Since the machine M_1 is deterministic, the machine M_2 has no invertible commands at all. The definition of the *symmetric* machine $M_3 = M_2^{sym}$ is the following. Suppose $M_2 = \langle X, Y, Q, \Theta, \vec{s}_1, \vec{s}_0\rangle$. Then by definition, $M_2^{sym} = \langle X, Y, Q, \Theta^{sym}, \vec{s}_1, \vec{s}_0\rangle$, where Θ^{sym} is the minimal *symmetric* set containing Θ, that is, with every command $[V_1 \to V'_1, \ldots, V_{k+1} \to V'_{k+1}]$ it contains the inverse command $[V'_1 \to V_1, \ldots, V'_{k+1} \to V_{k+1}]$; in other words, $\Theta^{sym} = \Theta^+ \sqcup \Theta^-$, where $\Theta^+ = \Theta$ and $\Theta^- = \{\theta^{-1} \mid \theta \in \Theta\}$.

[3] Here and in what follows we keep in mind that the difference $||w_i|| - |w_i|_a$ is a constant for any computation.

A computation $w_0 \to \cdots \to w_t$ of M_3 (or of another machine) is called *reduced* if its history is a reduced word. If the history $h \equiv \theta_1 \dots \theta_t$ contains a subword $\theta_i\theta_{i+1}$, where the commands θ_i and θ_{i+1} are mutually inverse, then obviously there is a shorter computation $w_0 \to \cdots \to w_{i-1} \equiv w_{i+1} \to \cdots \to w_t$ whose space does not exceed the space of the original computation.

Lemma 2.2. *Let $w \equiv w_0 \to w_1 \to \cdots \to w_t$ be an accepted reduced computation of the machine M_3, and the command θ_* be applicable to w. Then*

(a) the word $u(w)$ belongs to the language $\mathcal{L}$ recognized by M_1 and

(b) the space of this computation is at least $space_{M_1}(u(w))$.

Proof. Let $h \equiv \theta(1) \dots \theta(t)$ be the history of the computation. If for some i, $w_{i+1} = w_i \cdot \theta(i+1)$ where $\theta(i+1) \equiv \theta_{23}$ or $\theta(i+1)^{\pm 1}$ is a command of M_{23}, then one can modify our accepted computation so that, for $j > i+1$, every command $\theta(j)$ is a command of M_{23} and $||w_j|| \le ||w_{j-1}||$. Hence we may assume that h has exactly one letter θ_{23} followed by the commands of M_{23}, and $h \equiv h_0\tau_1 h_1 \tau_2 \dots \tau_s h_s$, where $\tau_s \equiv \theta_{23}$, $\tau_i \equiv \theta_{12}^{(-1)^{s-i-1}}$ for $i < s$, and the subwords h_i-s contain no connecting commands. We may also assume that the subword h_0 is empty since the command θ_* does not change the subword $u(w)$.

Since, by the $\vec{s}_{10}$-condition, only one command of the machine M_1 (and of its analog M_{22}) accepts, the last command of h_{s-1} is this unique command of M_{22}, and so this last command is positive. Therefore if h_{s-1} contains a letter θ^{-1}, where θ is a command of M_{22}, then h has a 2-letter subword $\theta_1^{-1}\theta_2$, where both θ_1 and θ_2 are commands of M_{22}. Hence there is a configuration w_i such that both θ_1 and θ_2 are applicable to w_i. This is impossible since the machine M_1 is deterministic and the history h_{s-1} is a reduced word. Therefore h_{s-1} is entirely the history of a computation of M_{22}, the computation $w_{i_{s-1}} \to \cdots \to w_t$ with history $\tau_{s-1}h_{s-1}\tau_s h_s \equiv \theta_{12}h_{s-1}\theta_{23}h_s$ is an accepting computation of M_2, the word $u(w_{i_{s-1}})$ belongs to the language $\mathcal{L}$, and the space of the computation $w_{i_{s-1}} \to \cdots \to w_t$ is at least $space_{M_1}(u(w_{i_{s-1}}))$ by Lemma 2.1 (c).

Assume, by induction on j, that $u(w_{i_{s-j}})$ belongs to $\mathcal{L}$ for $j \ge 1$, where the computation $w_{i_{s-j}} \to \cdots \to w_t$ has history $\tau_{s-j}h_{s-j} \dots \tau_s h_s$, and the space of this computation is at least $space_{M_1}(u(w_{i_{s-j}}))$.

Then the word $w_{i_{s-j-1}}$ has similar properties if h_{s-j-1} consists of the commands of M_{21} or their inverses since these commands do not change the content of the tapes numbered $1, \dots, k$. Otherwise the commands of h_{s-j-1} are commands of M_{22} (and their inverses), and since this machine is deterministic, the word h_{s-j-1} has no subwords $\theta_1^{-1}\theta_2$ with positive θ_1 and θ_2. Therefore we have $h_{s-j-1} \equiv g'g''^{-1}$, where both g' and g'' are (positive) histories of M_{22}-computations. This implies the equality $(w_{i_{s-j-1}} \cdot \tau_{s-j-1}) \cdot g' = w_{i_{s-j}} \cdot g''$. Since the commands $\theta_{12}^{\pm 1}$ do not change $u(w_i)$-s, we have $W_{i_{s-j-1}} \cdot g' = W_{i_{s-j}} \cdot g''$, where $W_{i_{s-j-1}}$ and $W_{i_{s-j}}$ are the input configurations for the machine M_1 with inputs $u(w_{i_{s-j-1}})$ and $u(w_{i_{s-j}})$, respectively. (Here we use identical letters for the corresponding commands of M_1 and M_{22}.)

The machine M_1 is deterministic, and so the accepted computation for $W_{i_{s-j}}$ must look like $W_{i_{s-j}} \to \cdots \to W_{i_{s-j}} \cdot g'' \to \dots$, and consequently, the configuration $W_{i_{s-j}} \cdot g''$ is accepted by M_1. Therefore we can construct the accepted computation $W_{i_{s-j-1}} \to \cdots \to W_{i_{s-j-1}} \cdot g' = W_{i_{s-j}} \cdot g'' \to \dots$ for M_1, and so the word $u(w_{i_{s-j-1}})$ belongs to $\mathcal{L}$, as desired.

The constructed accepted computation of M_1 is decomposed in two parts. It follows from the definition of M_{22} that the space of the first part is majorized by the space of the M_3-computation $w_{i_{s-j-1}} \to \cdots \to w_{i_{s-j-1}} \cdot g'$ which is a part of the computation $w_{i_{s-j-1}} \to \cdots \to w_t$. The second part occurs in a deterministic accepted M_1-computation with input $u(w_{i_{s-j}})$, and so, by the inductive hypothesis, the space of this part does not exceed the space of the M_3-computation $w_{i_{s-j}} \to \cdots \to w_t$. Hence $space_{M_1}(u(w_{i_{s-j-1}}))$ does not exceed the space of the M_3-computation $w_{i_{s-j-1}} \to \cdots \to w_t$.

Since $w \equiv w_{i_1}$, the lemma is proved by induction on j. □

Lemma 2.3. *The machines M_1 and M_3 recognize the same language. The generalized space complexities $S'_{M_2}(n)$ and $S'_{M_3}(n)$ are equivalent.*

Proof. We recall that every computation of the machine M_2 is also a computation of M_3. Therefore the first statement follows from Lemmas 2.1 (a) and 2.2 (a).

To prove the second part, it suffices to prove that for every accepted configuration w of M_2 (of M_3), there is an accepted configuration w' of M_3 (of M_2) such that $||w'|| \le ||w||$ but $space_{M_3}(w') \ge space_{M_2}(w)$ (respectively, $space_{M_2}(w') \ge space_{M_3}(w)$).

(1) Consider an accepted computation $w \equiv w_0 \to \cdots \to w_t$ of M_2 whose space is equal to $space_{M_2}(w)$. If the first command of this computation is not a command of M_{21}, then $||w_0|| \geq ||w_1|| \geq \cdots \geq ||w_t||$, and therefore $space_{M_2}(w) = |w|_a \leq space_{M_3}(w)$, so one can choose $w' \equiv w$. If the first command is a command of M_{21}, then by Lemmas 2.1(c) and 2.2 (b), we have $space_{M_2}(w) = \max(space_{M_1}(u(w)), |w|_a) \leq space_{M_3}(w)$, and again $w' \equiv w$.

(2) Now consider a reduced accepted computation $w \equiv w_0 \to \cdots \to w_t$ of M_3 whose space is equal to $space_{M_3}(w)$. If the first command (or its inverse) is a command of M_{23}, then the commands of the shortest accepted computation with minimal space just erase squares. Hence $space_{M_3}(w) = |w|_a = space_{M_2}(w)$, and we can choose w' equal to w. If the first command is a command of M_{21}^{sym}, then by Lemma 2.2 (a), the word w is accepted by M_2. Since every command of M_2 is a command of M_3, we have $space_{M_3}(w) \leq space_{M_2}(w)$, and again it suffices to set $w' \equiv w$.

Thus, we may assume that the first command of our computation (or its inverse) is a command of M_{22}. Therefore the history h of this computation has a prefix $h'h''$ with non-empty h'', where every command of h' is a command of M_{22}^{sym} and either every command of h'' (or the inverse command) is a command of M_{21} or every command of h'' is a command of M_{23}. In the latter case, we may assume that $h \equiv h'h''$ and $||w_0|| \geq ||w_1|| \geq \cdots \geq ||w_t||$, and so $space_{M_3}(w) = |w|_a \leq space_{M_2}(w)$, and $w' \equiv w$. In the former case, we set $w' = w \cdot h'$ and note that $||w|| = ||w_1|| = \cdots = ||w'||$ since the commands of the computation $w \to \cdots \to w'$ with history h' do not change the number of tape squares. In particular, we have $space_{M_3}(w) \leq space_{M_3}(w')$. Since the command θ_* is applicable to w' in this case, we have $space_{M_3}(w') \leq space_{M_2}(w')$ as was observed in the previous paragraph. Therefore $space_{M_3}(w) \leq space_{M_3}(w') \leq space_{M_2}(w')$, as desired; and the lemma is proved.

□

Lemma 2.4. *For every DTM M recognizing a language $\mathcal{L}$ and having space complexity $S(n)$ (for every NTM M recognizing a language $\mathcal{L}$ and having FSC space complexity $f(n)$), there exists an NTM M' with the following properties.*

1. *The machine M' recognizes the language $\mathcal{L}$.*
2. *M' is symmetric.*
3. *The space and the generalized space complexities of M' are equivalent to $S(n)$ (respectively, are equivalent to $f(n)^2$).*
4. *For every command $[V_1 \to V_1', \ldots, V_k \to V_k']$ of M', we have $\sum |V_i|_a + \sum |V_i'|_a \leq 1$, i.e., at most one tape letter is involved in the command.*

Proof. Assume that the machine M is deterministic. Then starting with $M_1 = M$, we construct the machine M_2 described in Subsection 2.2 and the machine $M_3 = M_2^{sym}$. Now Lemma 2.3 implies statement 1, and statement 2 is true since $M_3 = M_2^{sym}$. Then, by Lemmas 2.3 and 2.1 (b), we have

$$S'_{M_3}(n) \sim S'_{M_2}(n) \sim S_{M_2}(n) \sim S_{M_1}(n) = S(n),$$

and $S(n) \leq S_{M_3}(n) \leq S'_{M_3}(n)$ by Lemma 2.2 (b), and so all these functions are equivalent. Finally, we modify M_3 to obtain property 4. For example, if for a one-tape machine we have a command $aq \to bq'$, then we introduce a new state letter q'' and replace this command by two commands $aq \to q''$ and $q'' \to bq'$. It is easy to see that the obtained machine M' satisfies property 4 and inherits properties $1-3$ from M_3.

If M is non-deterministic, then we first use that the function $f(n)$ is FSC, and therefore, by Savitch's theorem ([9], Theorem 1.30), there exists a DTM M_1 accepting the same language L with space complexity equivalent to $f(n)^2$. So the replacement of $S(n)$ by $f(n)^2$ in the previous paragraph provides the proof of the non-deterministic version of the lemma. □

2.4 S-machines

By Lemma 3.9, computations of a machine are faithfully represented by special van Kampen diagrams (trapezia) over the group G. However this statement is true for S-machines but it is false for standard TM-s.

Ordinary Turing machines work with positive words and they can see some letters on the tape near the position where the head is. The command executed by the machine depends not only on the state of the head but also on the letter(s) observed by the head. In contrast, S-machines introduced in [SBR] work with words in group alphabets and they are almost "blind", i.e., the heads do not observe the tape letters. But the heads can "see" each other if there are no tape letters between them. We will use the following precise definition of an S-machine as a rewriting system.

Let k be a natural number. Consider a *language of admissible words*. It consists of words of the form

$$q_1 u_1 q_2 \dots u_k q_{k+1},$$

where q_i are letters from disjoint sets Q_i $(i = 1, \dots, k+1)$, u_i are reduced words in a group alphabet Y_i, (i.e. every letter a belongs to it iff the inverse letter a^{-1} does) and the sets $Y = \sqcup Y_i$ and $Q = \sqcup Q_i$ are finite. The letters from Q are called *state* letters, and the letters from Y are *tape* letters. Notice that in every admissible word, there is exactly one representative of each Q_i and these representatives appear in this word in the order of the indicees of Q_i. (i.e., unlike [24], we consider only the regular order of Q_i-s in admissible words).

There is a finite set of *commands* (or *rules*) Θ. To every $\theta \in \Theta$, we associate two sequences of reduced words from the free group $F(Q \cup Y)$: $B(\theta) = [U_1, ..., U_{k+1}]$, $T(\theta) = [V_1, ..., V_{k+1}]$, and a subset $Y(\theta) = \sqcup Y_i(\theta)$ of Y, where $Y_i(\theta) \subseteq Y_i$.

The words U_i, V_i satisfy the following restriction:

(*) For every $i = 1, ..., k+1$, the words U_i and V_i have the form

$$U_i \equiv v_{i-1} q_i u_i, \quad V_i \equiv v'_{i-1} q'_i u'_i$$

where $q_i, q'_i \in Q_i$, u_i and u'_i are words in the alphabet $Y_i(\theta)$, v_{i-1} and v'_{i-1} are words in the alphabet $Y_{i-1}(\theta)$. The words $v_0, v'_0, u_{k+1}, u'_{k+1}$ are empty.

Sometimes we will denote the rule θ by $[U_1 \to V_1, ..., U_{k+1} \to V_{k+1}]$. This notation contains no information about the sets $Y_i(\theta)$. In most cases it will be clear what these sets are. In the S-machines used in this paper, the sets $Y_i(\theta)$ will mostly be equal to either Y_i or $\emptyset$. By default $Y_i(\theta) = Y_i$.

We will use the notation $v_i q_i u_i \stackrel{\ell}{\to} v'_i q'_i u'_i$ for a part of a rule when the corresponding $Y_i(\theta)$ is empty (a similar notation has been used in [22]). In particular, u_i and u'_i are empty words in this case.

Every S-rule $\theta = [U_1 \to V_1, ..., U_{k+1} \to V_{k+1}]$ has an inverse $\theta^{-1} = [V_1 \to U_1, ..., V_{k+1} \to U_{k+1}]$; we set $Y_i(\theta^{-1}) = Y_i(\theta)$. We always divide the set of rules Θ of an S-machine into two disjoint parts, Θ^+ and Θ^- such that for every $\theta \in \Theta^+$, $\theta^{-1} \in \Theta^-$ and for every $\theta \in \Theta^-$, $\theta^{-1} \in \Theta^+$. The rules from Θ^+ (resp. Θ^-) are called *positive* (resp. *negative*).

To apply an S-rule θ to an admissible word $W \equiv q_1 w_1 q_2 \dots w_k q_{k+1}$ means to check if every w_i is a word in the alphabet $Y_i(\theta)$ and then, if W satisfies this condition, to simultaneously replace subwords U_i by subwords V_i $(i = 1, \dots, k+1)$. This replacement can be performed in the form $q_i \to v'_{i-1} v_i^{-1} q'_i u_i^{-1} u'_i$ followed by reducing the resulted word. The following convention is important in the definition of S-machine: *After every application of a rewriting rule, the word is automatically reduced. The reducing is not considered a separate step of an S-machine.*

The definitions of computation, its history, input admissible words, the accept word, the language of admissible words, space of a computation, space and generalized complexities, time and generalized time complexities of an S-machine are similar to those for a TM. (One should replace the word "configuration" by "admissible word" in the definitions.)

Although S-machines are usually highly non-deterministic, they are better adapted for simulating by finitely presented groups than ordinary TMs. On the other hand, it is mentioned in [27] that every symmetric NTM M can be viewed as an S-machine $S(M)$: just interpret the commands of the Turing machine as S-rules. (For example, the part of a rule of the form $aq \to bq'$ is interpreted as $q \to a^{-1} b q'$, and the part of the form $\alpha_j q \to \alpha_j q'$ is interpreted by the pair $\alpha_j \stackrel{\ell}{\to} \alpha_j$, $q \to q'$.) Unfortunately, the language recognized by $S(M)$ is in general much bigger than the language recognized by M since M works with a *positive* tape alphabet only. Nevertheless the following statement is true:

Lemma 2.5. *(Compare with Prop. 4.1[27].) Every computation of a symmetric NTM M is a computation of $S(M)$ with the same history. If M satisfies property 4 from Lemma 2.4, then every* positive *computation of $S(M)$, i.e., a computation consisting of positive words, is a computation of M with the same history.*

Proof. The first statement follows from the definition of the machine $S(M)$.

Every positive admissible word W of $S(M)$ is a configuration of the Turing machine M. Assume that a rule $\bar{\theta}$ of $S(M)$ corresponding to a command θ of M is applicable to this word W and the word $W \cdot \bar{\theta}$ is positive. Recall that by property 4 of Lemma 2.4, θ involves at most one tape letter (e.g., it cannot replace a tape letter by a tape letter or have a part of the form $aq \to aq'$). Therefore the positiveness of both W and $W \cdot \bar{\theta}$ implies that the application of $\bar{\theta}$ just coincides with the application of θ. The second statement of the lemma follows. □

2.5 Composition with an adding machine

One the one hand, S-machines much better fit for simulating their work by group relations than ordinary Turing machines (and moreover, S-machines are themselves treated in [24] as HNN-extensions of a free group with basis $Y \cup Q$). On the other hand, when a symmetric Turing mashine M starts working as an S-machine (see Subsection 2.4), the language of (positive) accepted words can enlarge uncontrolably. Therefore we will consider a composition $\mathcal{S}$ of a symmetric Turing machine M and an auxiliarly S-machine $Z(A)$ from [24]. Fortunately, this composition is an S-machine which inherits the language, the space function and the generalized space function of M.

In [24], the main duty of $Z(A)$ was the exponential slowing down of basic computations, while now we will mainly use the capacity of $Z(A)$ (observed in Lemma 3.25 (2) [24]) to check whether an admissible word is positive or not.

The tape alphabet of $Z(A)$ consists of an alphabet $A^{\pm 1}$ and two copies $A_1^{\pm 1}$ and $A_2^{\pm 1}$ of $A^{\pm 1}$ while the input alphabet is $A^{\pm 1}$. The admissible words are of the form $Lupv R$, where u is a reduced word in the alphabet $A^{\pm 1} \cup A_1^{\pm 1}$, v is a reduced word in $A_2^{\pm 1}$, the symbols L, p, R, are state letters, the commands do not change L and R, and $p \in \{p(1), p(2), p(3)\}$. The input words have the form $Lup(1)R$ and the accept ones are of the form $Lup(3)R$.

For every letter $a \in A$ let a_1, a_2 denote its copies in A_1, A_2.

Thus, the set of state letters is $Q_1 \cup Q_2 \cup Q_3$ where

$$Q_1 = \{L\}, Q_2 = \{p(1), p(2), p(3)\}, Q_3 = \{R\}.$$

The set of tape letters is $Y_1 \cup Y_2$ where $Y_1 = A \cup A_1$ and $Y_2 = A_2$.

The machine $Z(A)$ has the following positive rules (a is an arbitrary letter from A below). The following comments explain the meanings of these rules.

- $r_1(a) = [L \to L, p(1) \to a_1^{-1} p(1) a_2, R \to R]$.

  Comment. The state letter $p(1)$ moves left searching for a letter from A and replacing letters from A_1 by their copies in A_2.

- $r_{12}(a) = [L \to L, p(1) \to a^{-1} a_1 p(2), R \to R]$.

  Comment. When the first letter a of A is found, it is replaced by a_1, and $p(1)$ turns into $p(2)$.

- $r_2(a) = [L \to L, p(2) \to a p(2) a_2^{-1}, R \to R]$.

  Comment. The state letter $p(2)$ moves toward R.

- $r_{21} = [L \to L, p(2) \xrightarrow{\ell} p(1), R \to R]$, $Y_1(r_{21}) = Y_1, Y_2(r_{21}) = \emptyset$.

  Comment. $p(2)$ and R meet, and the cycle starts again.

- $r_{13} = [L \stackrel{\ell}{\to} L, p(1) \to p(3), R \to R]$, $Y_1(r_{13}) = \emptyset, Y_2(r_{13}) = A_2$.

  Comment. If $p(1)$ never finds a letter from A, then the cycle ends, and $p(1)$ turns into $p(3)$; p and L must stay next to each other in order for this rule to be executable.

- $r_3(a) = [L \to L, p(3) \to ap(3)a_2^{-1}, R \to R]$, $Y_1(r_3(a)) = A, Y_2(r_3(a)) = A_2$

  Comment. The letter $p(3)$ returns to R.

Here we will not explicitly use the form of these rules and will rather formulate, in Lemma 2.6, the required properties of $Z(A)$ obtained in [24]. Nevertheless the reader can see how this simple S-machine works if it 'agrees' to apply the rules in the order recommended in the comments. In particular, if the input word is $La^np(1)R$ $(n > 0)$ for some $a \in A$, then every maximal subcomputation ending with r_{21} and having no other rules r_{21} in its history, changes the word over Y_1 in the manner that one changes a binary n-digit number when one adds 1 to it. (Therefore the machine $Z(A)$ is called *adding*.) However Lemma 2.6 is not so obvious since (1) the machine $Z(A)$ is non-deterministic and (2) it works with arbitrary reduced tape words (while Turing machines deal with positive words only).

If w a word in the alphabet $A \cup A_1^{\pm 1} \cup A_2^{\pm 1}$, then its *projection* onto $A^{\pm 1}$ takes every letter to its copy in A.

Lemma 2.6. *The following properties of the machine $Z(A)$ hold.*

(1) Every positive input word u in the alphabet A is accepted by a canonical computation of $Z(A)$ with a positive history and such that all words appearing in this computation have equal lengths.

(2) For every computation $LupvR \equiv w \to \cdots \to w' \equiv Lu'p'v'R$ of $Z(A)$, the projections of the words uv and $u'v'$ onto A are freely equal. In particular, $u \equiv u'$ if the words v and v' are empty and the words u and u' contain no letters from $A_1^{\pm 1}$.

(3) If $w_0 \to \cdots \to w_t$ is a reduced computation of $Z(A)$ and $||w_0|| < ||w_1||$, then $||w_1|| \le ||w_2|| \le \cdots \le ||w_t||$.

(4) For every reduced computation $w_0 \to \cdots \to w_t$, we have $||w_i|| \le \max(||w_0||, ||w_t||)$ $(i = 0, \ldots, t)$.

(5) If $w \equiv LupR$, where $p \equiv p(1)$ (or $p \equiv p(3)$), $w \equiv w_0 \to \cdots \to w_t$ is a reduced computation, w_t contains the subword $p(3)R$ (respectively, $p(1)R$), and all a-letters of w_0 and w_t are from $A^{\pm 1}$, then u is a positive word and all words of the computation have the same length. The length of this computation is at least $2^{||u||}$.

(6) There is no reduced computation $w \equiv w_0 \to \cdots \to w_t$ of length $t \ge 1$ such that both w_0 and w_t contain $p(1)R$ or both of them contain $p(3)R$ and all a-letters of w_0 and w_t belong to $A^{\pm 1}$.

Proof. (1) To construct the canonical computation one should just use the above comments.

(2) This claim is the statement of Lemma 3.18 of [24].

(3) This statement is true by Lemma 3.24 of [24]

(4), (5) These statements are contained in Lemma 3.25 of [24].

(6) This statement is contained in Lemma 3.27 of [24].

□

Roughly speaking, the composition of machines M and $Z(A)$ was constructed in [24] as follows. After an application of (the copy of) a positive command from M, the copies of $Z(A)$ successivly implement their accepting computations on each of the tapes, and then (the copy of) the next command of M is applied. Thus $Z(A)$ just (exponentially) slows down the computations of M.

Here we somewhat modify the definition of the composition of a symmetric Turing machine M and the adding machine $Z(A)$. The difference is that machines of the form $Z(A)$ will work not only after the application of every command of M but before applications of commands from M as well. This makes it possible to simulate the work of any symmetric NTM, not only S-machines as was done in [24]. The aim of following construction is to obtain an S-machine $\mathcal{S}$ which recognizes the same language and has the same space and generalized space complexity as the symmetric Turing machine M. The S-machine constructed in [SBR] cannot serve in the present paper since the space and the generalized space complexities of that machine are equivalent to its time complexity.

Consider a symmetric NTM $M = \langle X, Y, Q, \Theta, \vec{s}_1, \vec{s}_0\rangle$ with $Y = \sqcup_{i=1}^{l} Y_i$, and with the $\vec{s}_{10}$-condition. The set Θ is a disjoint union of positive and negative commands: $\Theta = \Theta^+ \sqcup \Theta^-$. Let $S(M)$ be the associated S-machine defined before Lemma 2.5. We will assume that the admissible words of $S(M)$ are of the form $k_1u_1k_2 \dots u_lk_{l+1}$, where k_i are letters from disjoint sets Q_i $(i = 1, \dots, l+1)$, u_i is a reduced word in the alphabet Y_i $(i \le l)$.

To define the composition $\mathcal{S} = M \circ Z$ of M and $Z(A)$, we will insert a p-letter between any two consecutive q-letters k_i, k_{i+1} in an admissible word of $S(M)$, so that to treat any subword $k_i...p...k_{i+1}$ as an admissible word for a copy of $Z(A)$. In particular, the *start* and the *accept* words of $\mathcal{S}$ are obtained from, respectively, the start and accept words of M.

First, for every $i = 1, ..., l$, we make two copies of the alphabet Y_i of $S(M)$ $(i = 1, ..., l)$: $Y_{i,1}$ and $Y_{i,2}$. The set of state letters of the new machine is

$$K_1 \sqcup P_1 \sqcup K_2 \sqcup P_2 \sqcup ... \sqcup P_l \sqcup K_{l+1},$$

where $P_i = \{p_i, p_i(\theta, 1^-), p_i(\theta, 2^-), p_i(\theta, 3^-), p_i(\theta, 1^+), p_i(\theta, 2^+), p_i(\theta, 3^+) \mid \theta \in \Theta^+\}$, $i = 1, ..., l$.

The set of tape letters is

$$\bar{Y} = (Y_1 \sqcup Y_{1,1}) \sqcup Y_{1,2} \sqcup (Y_2 \sqcup Y_{2,1}) \sqcup Y_{2,2} \sqcup ... \sqcup (Y_l \sqcup Y_{l,1}) \sqcup Y_{l,2};$$

the components of this union will be denoted by $\bar{Y}_1, ..., \bar{Y}_{2l}$.

The set of positive rules $\bar{\Theta}$ of $M \circ Z$ is a union of the set of modified positive rules of $S(M)$ and of the positive rules of $Z_i(\theta, -)$ and $Z_i(\theta, +)$ $(\theta \in \Theta, i = 1, ..., l)$ which are copies of the machines $Z(Y_i)$ (also suitably modified).

More precisely, suppose a positive command θ of $S(M)$ differs from the unique start and accept commands of $S(M)$ and has the form

$$[k_1u_1 \to k_1'u_1', v_1k_2u_2 \to v_1'k_2'u_2', ..., v_lk_{l+1} \to v_l'k_{l+1}']$$

where $k_i, k_i' \in K_i$, u_i and v_i are words in Y_i. Then its copy in $M \circ Z$ is

$$\bar{\theta} = \begin{array}{l}[k_1u_1 \to k_1'u_1', v_1p_1(\theta, 3^-) \xrightarrow{\ell} v_1'p_1(\theta, 1^+), k_2u_2 \to k_2'u_2', ..., \\ v_lp_l(\theta, 3^-) \xrightarrow{\ell} v_l'p_l(\theta, 1^+), k_{l+1} \to k_{l+1}']\end{array}$$

with $\bar{Y}_{2i-1}(\bar{\theta}) = Y_i(\theta)$ and $\bar{Y}_{2i}(\theta) = \emptyset$ for every i, in particular, the words u_i, v_i are rewritten here in the alphabet Y_i.

Each machine $Z_i(\theta, -)$ is a copy of the machine $Z(Y_i)$, where every rule $\tau = [U_1 \to V_1, U_2 \to V_2, U_3 \to V_3]$ is replaced by rule of the form

$$\bar{\tau}_i(\theta, -) = \left[\begin{array}{l}\bar{U}_1 \to \bar{V}_1, \bar{U}_2 \to \bar{V}_2, \bar{U}_3 \to \bar{V}_3, \\ k_j \to k_j, p_j(\theta, 3^-) \xrightarrow{\ell} p_j(\theta, 3^-), j = 1, ..., i-1, \\ p_s(\theta, 1^-) \xrightarrow{\ell} p_s(\theta, 1^-), k_{s+1} \to k_{s+1}, s = i+1, ..., l\end{array}\right]$$

where $\bar{U}_1, \bar{U}_2, \bar{U}_3, \bar{V}_1, \bar{V}_2, \bar{V}_3$ are obtained from $U_1, U_2, U_3, V_1, V_2, V_3$, respectively, by replacing $p(j)$ with $p_i(\theta, j^-)$, L with k_i and R with k_{i+1}, and for $s \ne i$, $\bar{Y}_{2s-1}(\bar{\tau}_i(\theta, -)) = Y_i$.

Similarly, each machine $Z_i(\theta, +)$ is a copy of the machine $Z(Y_i)$, where every rule $\tau = [U_1 \to V_1, U_2 \to V_2, U_3 \to V_3]$ is replaced by the rule of the form

$$\bar{\tau}_i(\theta, +) = \left[\begin{array}{l}\bar{U}_1 \to \bar{V}_1, \bar{U}_2 \to \bar{V}_2, \bar{U}_3 \to \bar{V}_3, \\ k_j' \to k_j', p_j(\theta, 3^+) \xrightarrow{\ell} p_j(\theta, 3^+), j = 1, ..., i-1, \\ p_s(\theta, 1^+) \xrightarrow{\ell} p_s(\theta, 1^+), k_{s+1}' \to k_{s+1}', s = i+1, ..., l\end{array}\right]$$

where $\bar{U}_1, \bar{U}_2, \bar{U}_3, \bar{V}_1, \bar{V}_2, \bar{V}_3$ are obtained from $U_1, U_2, U_3, V_1, V_2, V_3$, respectively, by replacing $p(j)$ with $p_i(\theta, j^+)$, L with k_i' and R with k_{i+1}', and for $s \ne i$, $\bar{Y}_{2s-1}(\bar{\tau}_i(\theta, +)) = Y_i$.

If θ is the start (the accept) command of $S(M)$, then we introduce only machines $Z_i(\theta, +)$ (only $Z_i(\theta, -)$, respectively), and replace the letters $p_j(\theta, 3^-)$ (replace $p_j(\theta, 1^+)$, resp.) by p_j in the above definition of the command $\bar{\theta}$.

In addition, we need the following *transition* rules $\zeta(\theta,-)$ and $\zeta(\theta,+)$ that transform all p-letters from and to their original forms.

$$[k_i \to k_i, p_j \stackrel{\ell}{\to} p_j(\theta,1^-), i=1,...,l+1, j=1,...,l].$$

$$[k'_i \to k'_i, p_j(\theta,3^+) \stackrel{\ell}{\to} p_j, i=1,...,l+1, j=1,...,l].$$

If θ is the start (the accept) command of $S(M)$, then we introduce only $\zeta(\theta,+)$ (only $\zeta(\theta,-)$, respectively).

Thus while the machine $Z_i(\theta,-)$ (the machine $Z_i(\theta,+)$) works all other machines $Z_j(\theta,-)$ (all machines $Z_j(\theta,+)$, $j \neq i$) must stay idle (their state letters do not change and do not move away from the corresponding k-letters). After the machine $Z_i(\theta,-)$ (the machine $Z_i(\theta,+)$) finishes, i.e., the state letter $p_i(\theta,3^{\pm})$ appears next to k_{i+1} (next to k'_{i+1}), the next machine $Z_{i+1}(\theta,-)$ (the machine $Z_{i+1}(\theta,+)$) starts working. The transition rule $\zeta(\theta,-)$ switches on the consecutive works of the machines $Z_1(\theta,-),\dots,Z_l(\theta,-)$. After all p-letters have the form $p_j(\theta,3^-)$ we can apply the rule $\bar{\theta}$ and turn all $p_j(\theta,3^-)$ into $p_j(\theta,1^+)$. This switches on the consecutive work of $Z_1(\theta,+),\dots,Z_l(\theta,+)$, followed by the transition rule $\zeta(\theta,+)$.

So the modified rule $\bar{\theta} \in \bar{\Theta}$ turns on the machines $Z_1(\theta,+),\dots,Z_l(\theta,+)$ (and $\bar{\theta}$ follows after the machines $Z_1(\theta,-),\dots,Z_l(\theta,-)$ finish their work).

Thus, in order to simulate a computation of the symmetric TM M (and of the S-machine $S(M)$) consisting of a sequence of applications of rules $\theta_1,\theta_2,...,\theta_s$, we first apply all rules corresponding to θ_1 (before and after $\bar{\theta}$), then all rules corresponding to θ_2, then all rules corresponding to θ_3, etc. The language $\mathcal{L}_{\mathcal{S}}$ of $\mathcal{S}$ consists of some words u in the alphabet Y_1. In particular, every admissible input word of $\mathcal{S}$ is of the form $\Sigma(u) \equiv k_1 u p_1 k_2 p_2 k_3 \dots k_{l-1} p_l k_l$. We denote the accept word of $\mathcal{S}$ by $\Sigma_0 = \Sigma_0(\mathcal{S})$.

The modified rules $\bar{\theta}$ of $S(M)$ will be called *basic rules* of the S-machine $\mathcal{S}$.

2.6 Computations of machine $\mathcal{S}$

Lemma 2.10 will complete the part of our paper dealing with machines. This lemma makes possible to substitute the Turing machine M' from Lemma 2.4 by the S-machine $\mathcal{S}$. Lemma 2.9 (4) helps to obtain the lower bound for the space function of the group G in Subsection 5.2.

Given a computation C of $\mathcal{S}$, one obtains a computation $C_{S(M)}$ of $S(M)$ after omitting all the (copies of) commands of machines $Z_i(\theta,\pm)$ and deleting the additional state letters (letters of the sets $P_1,\dots,P_l$) from the admissible words of the computation C. (By Lemma 2.6 (2), the computation $C_{S(M)}$ is well-defined, although $C_{S(M)}$ is of length 0 if C has no basic rules.)

Lemma 2.7. *(1) The start and the accept rules of $\mathcal{S}$ are basic rules which are the copies of the start and the accept commands of M, respectively. The machine $M \circ Z$ satisfies the $\vec{s}_{10}$-condition.*

(2) If the history h of a reduced computation $C: \ w_0 \to \dots \to w_t$ of $\mathcal{S}$ contains no basic commands, then $||w_i|| \leq \max(||w_0||, ||w_t||)$ for every i $(0 \leq i \leq t)$. If the only basic letter of h is the last one, then $||w_i|| \leq ||w_0||$ for $i < t$.

(3) If a computation C of $\mathcal{S}$ is reduced then $C_{S(M)}$ is also reduced.

(4) For any reduced computation $C: \ w_0 \to \dots \to w_t$ of $\mathcal{S}$ starting and ending with basic rules we have $space_{\mathcal{S}}(C) = space_{S(M)}(C_{S(M)})$.

(5) For every positive reduced computation $C: \ w_0 \to \dots \to w_t$ of the machine $S(M)$, there is a canonical reduced computation $C_{\mathcal{S}}$ of $\mathcal{S}$ whose history starts and ends with basic commands, such that $(C_{\mathcal{S}})_{S(M)} = C$. Moreover, we have $space_{\mathcal{S}}(C_{\mathcal{S}}) = space_{S(M)}(C)$.

Proof. (1) Property (1) follows from the similar property of the machine M and from the definition of the machine $M \circ Z$.

(2) We start with the first statement. If the history of the whole computation consists of the commands of one machine $Z_i(\theta,\pm)$, then the statement follows from Lemma 2.6(3). Otherwise it has $s \geq 1$ admissible words $w_{i_1},\dots,w_{i_s}$, such that the history of every subcomputation

$$C_0 : w_0 \to \dots \to w_{i_1}, \dots, C_j : w_{i_j} \to \dots \to w_{i_{j+1}}, \dots, C_s : w_{i_s} \to \dots \to w_t$$

either consists of the commands of some $Z_i(\theta,\pm)$, being a maximal subcomputation with this property, or is equal to a transition letter $\zeta(\theta,\pm)^{\pm 1}$ for some θ. In the former case, all the admissible words participating in C_j have the same length for $j \in [1, s-1]$ by Lemma 2.6 (5,6). The same is clearly true in the latter case. Therefore it suffices to prove that the lengths of the admissible words do not decrease in the subcomputations C_0^{-1} and C_s.

Let us consider C_s only, assuming that it is a computation of some machine $Z_i(\theta,\pm)$. It corresponds to a computation $LupvR \equiv W_0 \to \cdots \to W_m \equiv Lu'pv'R$ of a machine of the form $Z(A)$ with $m = t - i_s$. Since $s \geq 1$ and C_s is a maximal subcomputation corresponding to $Z_i(\theta,\pm)$, we must have $||v|| = 0$. (Otherwise only commands of $Z_i(\theta,\pm)$ could be applied to w_{i_s}, and so $w_{i_s-1} \to w_{i_s} \to \cdots \to w_t$ is a longer computation of the same $Z_i(\theta,\pm)$.) Hence the projection of the word uv onto A is reducible, and so $||W_0|| \leq ||W_i||$ for every $i \in [1,m]$ by Lemma 2.6 (2). Therefore either all W_i-s have equal lengths or $||W_0|| \leq ||W_1|| \leq \cdots \leq ||W_m||$ by Lemma 2.6 (3). In any case, we have $||W_0|| \leq \cdots \leq ||W_m||$. This implies $||w_{i_s}|| \leq \cdots \leq ||w_t||$, as required.

The proof of the second claim is similar: The computation $w_{i_1} \to \cdots \to w_{i_s}$ is a product of subcomputations C_j-s which preserve the lengths of w_i-s, while C_0 and C_s cannot increase the space.

(3)Assume that $\tau_1 \dots \tau_t$ is a history of a computation $w_0 \to \cdots \to w_t$ of $\mathcal{S}$, where τ_1 corresponds to a positive command θ of $S(M)$, τ_t corresponds to θ^{-1}, and the other rules are not basic. Then the history h' of the computation $w_1 \to \cdots \to w_{t-1}$ is non-empty and $h' \equiv h_1 \dots h_s$, where each h_i $(i = 1, \dots, s)$ is a maximal subword of h' corresponding to the work of some machine $Z_{j(i)}(\theta_i,\pm)$. Since h_1 and h_s correspond to $Z_1(\theta,+)$, either $s = 1$ or there is i such that both h_{i-1} and h_{i+1} correspond to the same $Z_{j(i)\pm 1}(\theta',\pm)$. It follows that the computation with history h_i satisfies the assumption of Lemma 2.6 (6), a contradiction.

(4) Property (4) follows from (2) and the definition of the computation $C_{S(M)}$.

(5) Given C, the computation $C_{\mathcal{S}}$ with the same space and with property $(C_{\mathcal{S}})_{S(M)} = C$ is briefly described above at the end of subsection 2.5 and with more details (though with submachines $Z_i(\theta,+)$ but without $Z_i(\theta,-)$) in subsection 3.7 of [24]. □

Lemma 2.8. *Let $C:\ w_0 \to \cdots \to w_t$ be a reduced computation of $\mathcal{S}$ such that the first command $\bar\theta_1$ and the last commands $\bar\theta_t$ are basic ones, and $C_{S(M)} \equiv W_0 \to \cdots \to W_s$ with $s \geq 2$. Then the subcomputation $W_1 \to \cdots \to W_{s-1}$ is positive and $t - 2 \geq 2^{|W_1|_a/l}$. If $\bar\theta_1$ (if $\bar\theta_t$) is a start (is an accept) command, then the word W_0 (respectively, W_t) is also positive.*

Proof. To justify the first claim of the lemma, it suffices to prove that the word W_1 is positive under the assumption that there are no basic commands in the computation $w_1 \to \cdots \to w_{t-1}$, and the word W_1 corresponds to w_1. Therefore it suffices to prove that the word w_1 is positive.

We first assume that the first command $\bar\theta$ of the history of C is a positive basic command. Then $\bar\theta$ switches on the machine $Z_1(\theta,+)$. Since $s \geq 2$, this machine must complete its work before the computation C ends. The computation of $Z_1(\theta,+)$ cannot be empty since otherwise $\bar\theta$ would be followed by $\bar\theta^{-1}$ because w_1 involves the state letter $p_1(\theta,1^+)$. But this would contradict the reducibility of C.

Hence, by Lemma 2.6 (6), the work of $Z_1(\theta,+)$ sooner or later leads to an admissible word w_i containing the state letter $p_1(\theta,3^+)$. By Lemma 2.6(5), the subword of w_1 of the form $k_1u_1p_1(\theta,1^+)k_2$ is positive, and the time of the work of $Z_1(\theta,+)$ is at least $2^{||u_1||}$. Then the machine $Z_1(\theta,+)$ finishes working and switches on the machine $Z_2(\theta,+)$, whose work similarly provides the positiveness of the subword of w_1 having form $k_2u_2p_2(\theta,1+)k_3$. Finally we obtain that w_1 is covered by positive subwords, and so it is positive itself. The time of the work of all $Z_1(\theta,+), \dots, Z_l(\theta,+)$ is at least $2^{|W_1|_a/l}$ since $|W_1|_a = \sum_{j=1}^{l} ||u_j||$.

If the command $\bar\theta^{-1}$ is positive, then it switches on the machine $Z_l(\theta,-)$, and we first obtain the positiveness of $k_lu_lp_l(\theta,3^-)k_{l+1}$, then the positiveness of $k_{l-1}u_{l-1}p_{l-1}(\theta,3^-)k_l$, and so on.

Since start and accept commands leave tape letters unchanged, the second statement follows from the positiveness of the word W_1 (of the word W_{t-1}). The lemma is proved. □

Lemma 2.9. *(1) The S-machine $\mathcal{S}$ and the symmetric NTM M recognize the same language $\mathcal{L}$.*

(2) The space complexities $S_M(n)$ and $S_{\mathcal{S}}(n)$ of M and $\mathcal{S}$, respectively, are equivalent.

(3) The generalized space complexities $S'_M(n)$ and $S'_{\mathcal{S}}(n)$ of M and $\mathcal{S}$ are equivalent.

(4) We have $T'(n) \succeq \exp(S'_M(n))$ for the generalized time complexity $T'(n)$ of the machine $\mathcal{S}$.

Proof. (1) Assume that a word u belongs to the language $\mathcal{L}$ recognized by M. By Lemma 2.5, this word belongs to the language of $S(M)$, and the accepted computation C is positive. By Lemma 2.7(5), u belongs to the language $\mathcal{L}_{\mathcal{S}}$ of $\mathcal{S}$.

Now suppose u belongs to $\mathcal{L}_{\mathcal{S}}$, and C is the accepted computation. Then the computation $C_{S(M)}$ is positive by Lemma 2.8, and therefore this computation is also an accepted computation of the machine M by Lemma 2.5, and so $u \in \mathcal{L}$.

(2) The above argument shows that if a reduced computation $C : w_0 \to \cdots \to w_t$ of $\mathcal{S}$ accepts an admissible input word w_0, then the computation $C_{S(M)}$ is a positive accepting computation of both $S(M)$ and M, and so $S_M(n) \le S_{\mathcal{S}}(n)$ by Lemma 2.7 (4). On the other hand, every accepted input configuration W of M is accepted by $S(M)$. By Lemma 2.7(5), it has a copy accepted by $\mathcal{S}$, and moreover, the accepting computations of M and $\mathcal{S}$ need the same space. Therefore $S_M(n) \ge S_{\mathcal{S}}(n)$.

(3) Assume that $C : W \equiv W_0 \to \cdots \to W_t$ is an accepting computation of M such that $space_M(C) = space_M(W)$, and $C_{\mathcal{S}} : w \equiv w_0 \to \cdots \to w_s$. For the given w and w_s, we also consider a reduced computation $C' : w \to \cdots \to w_s$ of $\mathcal{S}$ with minimal space and the computation $(C')_{S(M)} : W'_0 \to \cdots \to W'_{t'}$. If $t' = 0$ (no basic rules), then $W_0 \equiv W_t$ by Lemma 2.6 (2), and so $space_M(C) = space_{\mathcal{S}}(C')$. Then we assume that $t' > 0$ and note that $||W'_0|| = ||W_0||$ and $||W'_{t'}|| = ||W_t||$ by Lemma 2.6 (2) since the corresponding admissible words of $\mathcal{S}$ can be connected by computations without basic rules. Since $C'_{S(M)}$ is positive by Lemma 2.8, it is also an accepted computation of the machine M by Lemma 2.5. Therefore, by Lemma 2.7 (4), $space_M(C) \le space_{S(M)}(C'_{S(M)}) = space_{\mathcal{S}}(C')$. Since $|w|_a = |W|_a$, the last inequality proves that $S'_M(n) \le S'_{\mathcal{S}}(n)$ for every n.

Now we consider any accepting computation $C : w \equiv w_0 \to \cdots \to w_t$ of $\mathcal{S}$ with $|w| \le n$, such that $space_{\mathcal{S}}(C) = space_{\mathcal{S}}(w)$. Without loss of generality, we may also assume that $space_{\mathcal{S}}(C[m]) = space_{\mathcal{S}}(w_m)$ for every subcomputation $C[m]$ of the form $w_m \to \cdots \to w_t$. Let $\rho_{i_1} \equiv \bar{\theta}_1, \ldots, \rho_t \equiv \rho_{i_s} \equiv \bar{\theta}_s$ be the basic commands of the history $h \equiv \rho_1 \ldots \rho_t$ of C, let C' be the subcomputation of C with history $h' \equiv \rho_1 \ldots \rho_{i_1 - 1}$, and let C'' have history $h'' \equiv \rho_{i_1} \ldots \rho_t$; and so $h \equiv h'h''$ and $C = C'C''$.

We denote by $W_0 \to W_1 \to \cdots \to W_s$ the computation $(C'')_{S(M)}$ and by $(C'')_{S(M)}[1]$ the subcomputation $W_1 \to \cdots \to W_s$, which is positive by Lemma 2.8. Note that $space_{S(M)}((C'')_{S(M)}[1]) = space_{S(M)}(W_1)$ since otherwise the subcomputation of $C[i_1]$ could be replaced by a subcomputation which needs less space by Lemma 2.7 (5). Therefore by Lemma 2.7 (4),

$$space_{\mathcal{S}}(C'') = space_{S(M)}((C'')_{S(M)}) \le$$

$$space_{S(M)}((C'')_{S(M)}[1]) + |||W_1|| - ||W_0||| = space_{S(M)}(W_1) + c$$

since $|||W_1|| - ||W_0|||$ is bounded by a constant c depending on the machine M only.

By Lemma 2.7 (2), we have $||w_j|| \le ||w_0||$ for $j \le i_1$, and so $space_{\mathcal{S}}(C') \le |w_0|_a$. Now, since $||W_1|| \le ||W_0|| + c \le ||w_{i_1 - 1}|| + c \le ||w_0|| + c$, we get

$$space_{\mathcal{S}}(C) \le \max(space_{\mathcal{S}}(C'), space_{\mathcal{S}}(C'')) \le$$

$$\max(|w_0|_a, space_{S(M)}(W_1) + c) \le \max(|w_0|_a, S'_{S(M)}(|w_0|_a + c) + c)$$

Hence $S'_{\mathcal{S}}(n) \le \max(S'_{S(M)}(n + c) + c, n)$. This inequality together with the inequality $S'_M(n) \le S'_{\mathcal{S}}(n)$ obtained earlier, show that $S'_M(n) \sim S'_{\mathcal{S}}(n)$.

(4) Let again $C : W \equiv W_0 \to \cdots \to W_t$ be an accepting computation of M such that $space_M(C) = space_M(W) = S'_M(|W|_a)$, and $C_{\mathcal{S}} : w \equiv w_0 \to \cdots \to w_s$. For the given w and w_s, we also consider a reduced computation $C' : w \to \cdots \to w_{s'}$ of $\mathcal{S}$ with minimal *time* s' and the computation $(C')_{S(M)} : W'_0 \to \cdots \to W'_{t'}$, where one has $W'_0 \equiv W_0$ and $W'_{t'} \equiv W_t$. Therefore $(C')_{S(M)}$ is a positive computation by Lemma 2.8, and by the choice of C and by Lemma 2.5, $space_M(C) \le space_M((C')_{S(M)})$. Consider also a maximal subcomputation C'' of C' starting and ending with basic commands. Then $(C')_{S(M)} = (C'')_{S(M)}$. By Lemma 2.8, $space_{S(M)}(C'')_{S(M)} \le c \log(time_{\mathcal{S}}(C''))$ for a constant $c > 0$. Thus by Lemma 2.5,

$$S'_M(|W|_a) = space_M(C) \le space_M((C')_{S(M)}) = space_{S(M)}((C')_{S(M)})$$

$$= space_{S(M)}((C'')_{S(M)}) \le c \log(time_{\mathcal{S}}(C'')) \le c \log(time_{\mathcal{S}}(C')) \le c \log(T'(|W|_a)),$$

and the lemma is proved.

□

Lemma 2.10. *(a) For every DTM M recognizing a language $\mathcal{L}$ and having a space complexity $S(n)$, (b) for every NTM M recognizing a language $\mathcal{L}$ and having an FSC space complexity $f(n)$, there exists an S-machine $\mathcal{S}$ with the following properties.*

1. *The machine $\mathcal{S}$ recognizes the language $\mathcal{L}$.*
2. *Respectively, (a) both the space and the generalized space complexities of $\mathcal{S}$ are equivalent to $S(n)$, (b) both the space and the generalized space complexities of $\mathcal{S}$ are equivalent to $f(n)^2$.*
3. *Every command of $\mathcal{S}$ or its inverse inserts/deletes at most one letter on the left and at most one letter on the right of every state letter.*
4. *The machine $\mathcal{S}$ satisfies the $\vec{s}_{10}$-property.*
5. *The unique start command is of the form $q_1 \to q'_1, q_2 \stackrel{\ell}{\to} q'_2, \ldots, q_{k+1} \stackrel{\ell}{\to} q'_{k+1}$, where $(q_1, \ldots, q_{k+1}) \equiv \vec{s}_1$.*
6. *Any state letters q from Q_1 is passive, i.e., there are no commands of $\mathcal{S}$ of the form $q \to q'u$ with non-empty a-word u.*

Proof. For every DTM M recognizing a language $\mathcal{L}$ and having a space complexity $S(n)$ (for every NTM M recognizing a language $\mathcal{L}$ and having an FSC space complexity $f(n)$) one can construct a symmetric NTM M', as in the formulation of Lemma 2.4. Then the properties 1 and 2 hold for the machine $\mathcal{S} = M' \circ Z$ by Lemma 2.9.

To provide the third property we use the same trick as for property 4 of Lemma 2.4. The properties 1 and 2 are obviously preserved.

To obtain the $\vec{s}_{10}$-condition of $\mathcal{S}$, it suffices to add new state letters and two new (positive) commands. Such a modification preserves the other properties of $\mathcal{S}$, as this was noticed at the end of subsection 2.1. Now the $\vec{s}_{10}$-condition for $\mathcal{S}$ follows from Lemma 2.7 (1). The form of the start command follows from our agreement that all tapes numbered $2, \ldots, k$ are empty for the input configurations of the machine M. Finally, the left-most head of the Turing machine M is passive being equal to the separating symbol α_1; and the same property is inherited by $\mathcal{S}$ as this follows from the definition of the composition $M' \circ Z$. □

3 Groups and diagrams

Recall that given a finitely generated group H, we want to construct an embedding of H into a finitely presented group G whose space function is equivalent to the deterministic space complexity of the algorithmic word problem in H. The relations of G are presented in this section.

3.1 Construction of the embedding

Let H be a finitely generated group with solvable word problem. To prove Theorem 1.2, we will suppose that a Turing machine M solves the word problem in H. This implies that H has a finite set of generators $\{a_1, .., a_m\}$, and a word w in these generators is accepted by M iff $w = 1$ in H. Here we consider only positive words in the generators since M can work with positive words only, and so we assume that the set of generators is symmetric: for every a_i, there is a generator a_j such that $a_i a_j = 1$ in H. Thus every relation holding in H follows from relations in $a_1, \ldots, a_m$, with positive left-hand side.

Further we will assume that one of the hypotheses (a), (b) of Lemma 2.10 holds. Therefore we also have the S-machine $\mathcal{S} = \langle X, Y, Q, \Theta, \vec{s}_1, \vec{s}_0 \rangle$ provided by that lemma. The input alphabet of $\mathcal{S}$ is the system of generators $a_1, \ldots, a_m$ of the group H together with the symbols of the inverse letters $\{a_1^{-1}, .., a_m^{-1}\}$. Let $S'_{\mathcal{S}}(n)$ be the generalized space complexity of $\mathcal{S}$.

We denote by $\hat{\mathcal{S}} = \langle \hat{X}, \hat{Y}, \hat{Q}, \hat{\Theta}, \hat{\vec{s}}_1, \hat{\vec{s}}_0 \rangle$ a copy of the S-machine $\mathcal{S}$. We will assume that $\hat{X} = X$, $Y \cap \hat{Y} = X$, $Q \cap \hat{Q}$ consists of the state letters of the vectors $\vec{s}_1$, and $\vec{s}_0$, and $\Theta \cap \hat{\Theta} = \emptyset$. Therefore the machines $\mathcal{S}$ and $\hat{\mathcal{S}}$ have the same admissible input words.

The copy of a command θ of $\mathcal{S}$ is called $\hat{\theta}$. Similar notation is used for the a-letters from $\hat{Y}$ and q-letters from $\hat{Q}$. The set of rules of the machine and admissible words of $\mathcal{S} \cup \hat{\mathcal{S}}$ is, by definition, the union of the corresponding sets for $\mathcal{S}$ and for $\hat{\mathcal{S}}$. The following lemma is a clear consequence of these definitions.

Lemma 3.1. *The sets of accepted admissible input words of the machines $\mathcal{S}$, $\hat{\mathcal{S}}$ and $\mathcal{S} \cup \hat{\mathcal{S}}$ coincide, and so these machines recognize the same language $\mathcal{L}$. They also have equal space complexities and equal generalized space complexities. For every accepting computation $w_1 \to \dots$ of $\mathcal{S} \cup \hat{\mathcal{S}}$, there is an accepting computation $w_1 \to \dots$ of either $\mathcal{S}$ or $\hat{\mathcal{S}}$ whose length and space does not exceed the length and space of the original computation.*

We consider a group $G(\mathcal{S}, L)$ associated with the machine $\mathcal{S}$. Furthermore as in [23], we need a very similar group $\hat{G}(\mathcal{S}, L)$ to produce a group embedding required for the proof of Theorem 1.2.

To define $G(\mathcal{S}, L)$ we need many copies of every letter used in the work of $\mathcal{S}$; this enables us to apply a kind of hyperbolic argument for the hub structure of van Kampen diagrams. Moreover, the copies alternate with "mirror copies"; this trick will be used in Section 4.

Therefore we "multiply" the machine $\mathcal{S}$ as follows. For some even $L \geq 40$, we introduce $L/2$ copies $\mathcal{S} = \mathcal{S}_1, \mathcal{S}_3, \dots, \mathcal{S}_{L-1}$ of the machine $\mathcal{S}$ and $L/2$ mirror copies $\mathcal{S}_2, \mathcal{S}_4, \dots, \mathcal{S}_L$ of $\mathcal{S}$. (A mirror copy of an arbitrary word $x_1 \dots x_n$ is, by definition, $x_n \dots x_1$. For even i, the rules of $\mathcal{S}_i(L)$ transform the words in the mirror manner in comparison with $\mathcal{S}$.) We also add auxiliary separating state letters $k_1, \dots k_L$. For every admissible word W of the machine $\mathcal{S}$, we define the words $W_1, W_2, \dots, W_L$, where $W_1, W_3, \dots W_{L-1}$ are copies of W in disjoint alphabets and $W_2, W_4, ..., W_L$ are mirror copies of W also in disjoint alphabets. The words of the form $k_1 W_1 k_2 W_2 \dots k_L W_L$ are admissible words of the machine $\mathcal{S}(L)$. The rules of $\mathcal{S}(L)$ are in one-to-one correspondence with the rules of $\mathcal{S}$; they transform the words $W_1, W_3, \dots$ as the rules of $\mathcal{S}$ do, they transform the words $W_2, W_4, \dots$ in the mirror manner, and do not change the new state letters $k_1, \dots, k_L$. We identify the set of rules of $\mathcal{S}(L)$ with Θ. Thus, by definition of $\mathcal{S}(L)$, we also multiply the input and accept configurations. If u is an input word for $\mathcal{S}$, then we denote by $\Sigma(u, L)$ the corresponding input configuration for $\mathcal{S}(L)$ containing $L/2$ copies of u and $L/2$ mirror copies of u as subwords. If the admissible words of $\mathcal{S}$ have K state letters, then the admissible words of the machine $\mathcal{S}(L)$ have $N = (K+1)L$ state letters. Clearly the machine $\mathcal{S}(L)$ enjoys the properties (1)–(5) of machine $\mathcal{S}$ listed in Lemma 2.10.

The finite set of generators of the group $G(\mathcal{S}, L)$ consists of *q-letters* corresponding to the states of $\mathcal{S}(L)$, *a-letters* corresponding to the tape letters of $\mathcal{S}(L)$, and *θ-letters* corresponding to the commands. Thus the set of generators consists of the set of (state) q-letters $Q(\mathcal{S}, L) = \sqcup_{i=1}^{N} Q_i$, the set of (tape) a-letters $Y = \sqcup_{i=1}^{N} Y_i$ including the input alphabet $X = \sqcup_{i=1}^{L} X_i$; and the θ-letters from N copies of Θ^+, i.e., for every $\theta \in \Theta^+$, we have N generators $\theta_1, \dots, \theta_N$.

The relations of the group $G(\mathcal{S}, L)$ correspond to the rules of the machine $\mathcal{S}(L)$; for every $\theta = [U_1 \to V_1, \dots U_N \to V_N] \in \Theta^+$, we have

$$U_i \theta_{i+1} = \theta_i V_i, \qquad \theta_j a = a \theta_j, \quad i, j = 1, ..., N \tag{3.4}$$

for all $a \in \bar{Y}_j(\theta)$. (Here $\theta_{N+1} \equiv \theta_1$.) The first type of relations will be called (θ, q)-*relations*, the second type - (θ, a)-*relations*.

The definition of the machine $\hat{S}(L)$ is similar to that of $\mathcal{S}(L)$ but the admissible words are of the form $k_1 \hat{W}_1 k_2 \hat{W}_2 ... k_L \hat{W}_L$, where every $\hat{W}_i$ is obtained from W_i after replacement of every letter x by its copy $\hat{x}$, and for $i = 1$, we, in addition, delete all a-letters, i.e., the word $\hat{W}_1$ has no a-letters. (In other words, instead of the first copy of $\mathcal{S}$, we use the "machine" with the same state letters but having no tape letters.) In particular, the word $\hat{\Sigma}(u, L)$ is obtained from $\Sigma(u, L)$ by omitting the first occurrence of the input word u. Again, it is obvious that properties (1) - (5) from Lemma 2.10 hold for the machine $\hat{\mathcal{S}}(L)$ as well. The relations of the group $\hat{G}(\mathcal{S}, L)$ are

$$\hat{U}_i \hat{\theta}_{i+1} = \hat{\theta}_i \hat{V}_i, \quad i = 1, ..., N, \qquad \hat{\theta}_j \hat{a} = \hat{a} \hat{\theta}_j \tag{3.5}$$

for all $\hat{a} \in Y_j(\hat{\theta})$ and $j \in [K+1, N]$.

We will also use the combined S-machine $S(L) \cup \hat{\mathcal{S}}(L)$. Its admissible words are either the admissible words for $\mathcal{S}(L)$ or the admissible words for $\hat{\mathcal{S}}(L)$, the set of rules is the union of the rules for $\mathcal{S}(L)$ and $\hat{\mathcal{S}}(L)$. Note that the machine $S(L) \cup \hat{\mathcal{S}}(L)$ does not satisfy the $\vec{s}_{10}$-condition.

The subwords of the form $(k_i W_i k_{i+1})^{\pm 1}$ (indices taken modulo L) of the admissible words of the machine $\mathcal{S}(L)$ are said to be i-sector words. Similarly one define i-sector words for the machines $\hat{\mathcal{S}}(L)$ and for $\mathcal{S}(L) \cup \hat{\mathcal{S}}(L)$. (Recall that the q-letters of the 1-sector words of $\mathcal{S}(L)$, except for k_1 and k_2, are identified with the corresponding letters of the original S-machine $\mathcal{S}$.) The state letters of the i-sector of $\mathcal{S}(L)$ are the letters from $\sqcup_{j=(i-1)(K+1)+1}^{i(K+1)} Q_j$, and the tape letters of the i-sector are the letters from $\sqcup_{j=(i-1)(K+1)+1}^{i(K+1)} Y_j$. Similarly we have state and tape letters of the i-sector for $\hat{\mathcal{S}}(L)$ and for $\mathcal{S}(L) \cup \hat{\mathcal{S}}(L)$. By definition, the θ-letters θ_j and $\hat{\theta}_j$ with subscripts $j \in [(i-1)(K+1)+1, i(K+1)]$ are θ-letters of the i-sector of $\mathcal{S}(L) \cup \hat{\mathcal{S}}(L)$. The state, tape and the θ-letters of the i-sector constitute the *alphabet* $\mathcal{A}_i$ of the i-sector.

The group $G(\mathcal{S} \cup \hat{\mathcal{S}}, L)$ is given by the generators and relations of both groups $G(\mathcal{S}, L)$ and $\hat{G}(\mathcal{S}, L)$.

Finally, the required group G is given by the generators and relations of the group $G(\mathcal{S} \cup \hat{\mathcal{S}}, L)$ and one more additional relation, namely the *hub*-relation

$$\Sigma_0 = 1, \tag{3.6}$$

where $\Sigma_0 = \Sigma(L)$ is the accept word (of length N) of the machine $\mathcal{S}(L)$ (and of $\hat{\mathcal{S}}(L)$ as well).

Suppose an admissible word W' of $\mathcal{S}(L)$ is obtained from an admissible word W by an application of a rule $\theta : [U_1 \to V_1, \dots, U_N \to V_N]$. This implies that $W = U_1 w_1 U_2 w_2 \dots U_N w_N$, where w_i is a word in the alphabet $Y_i(\theta)$, and therefore $W' = \theta_1^{-1} W \theta_1$ in $G(\mathcal{S}, L)$ by Relations (3.4), since $\theta_1 \equiv \theta_{N+1}$.

Now suppose u is a positive word in the alphabet $\{a_1, \dots, a_m\}$ vanishing in the group H. Then it is recognized by the machine $\mathcal{S}$, and so the word $\Sigma(u)$ is accepted by the machine $\mathcal{S}(L)$ and therefore it is conjugate to Σ_0 in the group $G(\mathcal{S} \cup \hat{\mathcal{S}}, L)$. Consequently, we have that $\Sigma(u) = 1$ in G by (3.6). Similarly, $\hat{\Sigma}(u) = 1$ in G.

Recall that we have identified the alphabet of 1-sector words of $\mathcal{S}(L)$ with the alphabet of $\mathcal{S}$. Therefore the word $\hat{\Sigma}(u)$ results from $\Sigma(u)$ after deleting the subword u in the alphabet of generators of H. Hence the relations $\Sigma(u) = \hat{\Sigma}(u) = 1$ imply $u = 1$ in G. Since the language of accepted words for $\mathcal{S}(L)$ contains all the defining relations of H, we have obtained

Lemma 3.2. *The mapping $a_i \mapsto a_i$ $(i = 1, \dots, m)$ extends to a homomorphism of the group H to G.*

In Section 4 we show that this homomorphism is injective.

3.2 Minimal diagrams

As in [21], we enlarge the set of defining relations of the group G by adding some consequences of defining relations. Taking into account Lemma 3.2, we include all cyclically reduced relations of the group H generated by the set $\{a_1, \dots, a_m\}$, i.e., all non-empty cyclically reduced words in $\{a_1^{\pm 1}, \dots, a_m^{\pm 1}\}$ which are equal to 1 in the group H. These relations will be called *H-relations*.

We denote by G_1 the group given by all generators of the group G, by all H-relations, and by all defining relations of G except for the hub-relation (3.6).

Recall that a van Kampen *diagram* Δ over a presentation $P = \langle B \mid \mathcal{R} \rangle$ (or just over the group P) is a finite oriented connected and simply–connected planar 2–complex endowed with a labeling function $\phi : E(\Delta) \to B^{\pm 1}$, where $E(\Delta)$ denotes the set of oriented edges of Δ, such that $\phi(e^{-1}) \equiv \phi(e)^{-1}$. Given a cell (that is a 2-cell) Π of Δ, we denote by $\partial\Pi$ the boundary of Π; similarly, $\partial\Delta$ denotes the boundary of Δ. The labels of $\partial\Pi$ and $\partial\Delta$ are defined up to cyclic permutations. An additional requirement is that the label of any cell Π of Δ is equal to (a cyclic permutation of) a word $R^{\pm 1}$, where $R \in \mathcal{R}$. Labels and lengths of paths are defined as for Cayley graphs.

The van Kampen Lemma states that a word W over the alphabet $B^{\pm 1}$ represents the identity in the group P if and only if there exists a diagram Δ over P such that $\phi(\partial\Delta) \equiv W$ ([17], Ch. 5, Theorem 1.1).

We will study diagrams over the groups G and G_1. The edges labeled by state letters ($= q$-*letters*) will be called *q-edges*, the edges labeled by tape letters (= *a-letters*) will be called *a-edges*, and the edges labeled by the letters from Θ and $\hat{\Theta}$ (= *θ-letters*) are *θ-edges*. The cells corresponding to Relation (3.6) are called *hubs*, the cells corresponding to Relations (3.4) and (3.5) are called *(θ, q)-cells* if they involve

q-letters, and they are called (θ, a)-*cells* otherwise. The cells corresponding to arbitrary relations of H are H-*cells.*

The resulting presentation and the diagrams over it are *graded* by the ranks of defining words and cells as follows. The hubs are the cells of the highest rank, the rank of (θ, q)-cells is higher than the rank of H-cells (and in Lemma 5.1, (θ, k_i)-cells, with $i \neq 1, 2$ are higher than other (θ, q)-cells), and the (θ, a)-cells are of the lowest rank.

If Δ and Δ' are diagrams over G, then we say that Δ has a higher type than Δ' if Δ has more hubs, or if the numbers of hubs are the same, but Δ has more cells which are next in the hierarchy, and so on.

Clearly the defined partial order on the set of diagrams satisfies the descending chain condition, and so there is a diagram having the smallest type among all diagrams with the same boundary label. Such a diagram is called *minimal.*

3.3 Bands and trapezia

From now on, we shall mainly consider minimal van Kampen diagrams. In particular the diagrams are *reduced*, i.e., they contain no pair of cells that have a common edge e such that the boundary paths of these cells starting with e have the same label. As in [27], [4], to explore van Kampen diagrams over the groups G and G_1 we shall use their simpler subdiagrams such as bands and trapezia.

Here we repeat some necessary definitions.

Definition 3.3. Let $\mathcal{Z}$ be a subset of the set of generators $\mathcal{X}$ of the group G. A $\mathcal{Z}$-band $\mathcal{B}$ is a sequence of cells $\pi_1, ..., \pi_n$ in a van Kampen diagram such that

- Every two consecutive cells π_i and π_{i+1} in this sequence have a common edge e_i labeled by a letter from $\mathcal{Z}$.
- Each cell π_i, $i = 1, ..., n$ has exactly two $\mathcal{Z}$-edges, e_{i-1} and e_i (i.e. edges labeled by a letter from $\mathcal{Z}$).
- If $n = 0$, then $\mathcal{B}$ is just a $\mathcal{Z}$-edge.

The counterclockwise boundary of the subdiagram formed by the cells $\pi_1, ..., \pi_n$ of $\mathcal{B}$ has the factorization $e^{-1}q_1 f q_2^{-1}$ where $e = e_0$ is a $\mathcal{Z}$-edge of π_1 and $f = e_n$ is an $\mathcal{Z}$-edge of π_n. We call q_1 the *bottom* of $\mathcal{B}$ and q_2 the *top* of $\mathcal{B}$, denoted $\mathbf{bot}(\mathcal{B})$ and $\mathbf{top}(\mathcal{B})$. Top/bottom paths and their inverses are also called the *sides* of the band. The $\mathcal{Z}$-edges e and f are called the *start* and *end* edges of the band. If $n \geq 1$ but $e = f$, then the $\mathcal{Z}$-band is called a $\mathcal{Z}$-*annulus.*

We say that a $\mathcal{Z}_1$-band and a $\mathcal{Z}_2$-band *cross* if they have a common cell and $\mathcal{Z}_1 \cap \mathcal{Z}_2 = \emptyset$.

We shall call a $\mathcal{Z}$-band *maximal* if it is not contained in any other $\mathcal{Z}$-band.

We will consider q-bands where $\mathcal{Z}$ is one of the sets Q_i of state letters for the machine $\mathcal{S}(L) \cup \hat{\mathcal{S}}(L)$, θ-bands for every $\theta \in \Theta$, and a-bands where $M = \{a\} \subseteq Y$.

The convention is that a-bands do not contain q-cells, and so they consist of (θ, a)-cells only.

The papers [20], [4], [23] contain the proof of the following lemma in a more general setting. (In contrast to Lemmas 6.1 [20] and 3.11 [23], we have no x-cells here.)

Lemma 3.4. *A minimal van Kampen diagram Δ over G_1 has no q-annuli, no θ-annuli, and no a-annuli. Every θ-band of Δ shares at most one cell with any q-band and with any a-band.*

If $W \equiv x_1 ... x_n$ is a word in an alphabet X, Y is another alphabet, and $\phi\colon X \to Y \cup \{1\}$ (where 1 is the empty word) is a map, then $\phi(W) \equiv \phi(x_1) ... \phi(x_n)$ is called the *projection* of W onto Y. We shall consider the projections of words in the generators of G onto $\Theta \sqcup \hat{\Theta}$ (all θ-letters map to the corresponding element of $\Theta \sqcup \hat{\Theta}$, all other letters map to 1), and the projection onto the alphabet $\{Q_1 \sqcup \cdots \sqcup Q_N\}$ (every q-letter maps to the corresponding Q_i, all other letters map to 1).


Definition 3.5. The projection of the label of a side of a q-band onto the alphabet $\Theta^{\pm 1}$ is called the *history* of the band. The projection of the label of a side of a θ-band onto the alphabet $\{Q_1, ..., Q_n\}$ is called the *base* of the band. Similarly we define the history of a word and the base of a word. The base of a word W is denoted by $base(W)$. It will be convenient to use representatives of $Q_1, ..., Q_N$ in base words. For example, if $k \in Q_1$, $q \in Q_2$, we shall say that the word kaq has base kq instead of $Q_1 Q_2$.

Definition 3.6. Let Δ be a minimal van Kampen diagram over G_1 which has boundary path of the form $p_1^{-1}q_1p_2q_2^{-1}$ where:

(TR_1) p_1 and p_2 are sides of q-bands,

(TR_2) q_1, q_2 are maximal parts of the sides of θ-bands such that $\phi(q_1)$, $\phi(q_2)$ start and end with q-letters,

(TR_3) for every θ-band $\mathcal{T}$ in Δ, the labels of $\mathbf{top}(\mathcal{T})$ and $\mathbf{bot}(\mathcal{T})$ are reduced.



Then Δ is called a *trapezium*. The path q_1 is called the *bottom*, the path q_2 is called the *top* of the trapezium, the paths p_1 and p_2 are called the *left and right sides* of the trapezium. The history of the q-band whose side is p_2 is called the *history* of the trapezium; the length of the history is called the *height* of the trapezium. The base of q_1 is called the *base* of the trapezium.

Remark 3.7. (1) Property (TR_3) is easy to achieve: by folding edges with the same labels having the same initial vertex, one can make the boundary label of a subdiagram in a van Kampen diagram reduced, see [27].

(2) Notice that the top (bottom) side of a θ-band $\mathcal{T}$ does not necessarily coincide with the top (bottom) side q_2 (side q_1) of the corresponding trapezium of height 1, and q_2 (q_1) is obtained from $\mathbf{top}(\mathcal{T})$ (resp. $\mathbf{bot}(\mathcal{T})$) by trimming the first and the last a-edges if these paths start and/or end with a-edges. We shall denote the trimmed top and bottom sides of $\mathcal{T}$ by $\mathbf{ttop}(\mathcal{T})$ and $\mathbf{tbot}(\mathcal{T})$. By definition, for arbitrary θ-band $\mathcal{T}$, $\mathbf{ttop}(\mathcal{T})$ is obtained by such a trimming only if $\mathcal{T}$ starts and/or ends with a (θ, q)-cell; otherwise $\mathbf{ttop}(\mathcal{T}) \equiv \mathbf{top}(\mathcal{T})$. The definition of $\mathbf{tbot}(\mathcal{T})$ is similar.

The trapezium Δ is said to be an i-sector if the labels of its top and bottom paths are i-sector words.

Lemma 3.8. *Let Γ be an i-sector, where $i \neq 1$. Then the sides p_1 and p_2 are the sides of maximal k_i- and k_{i+1}-bands $\mathcal{K}_i$ and $\mathcal{K}_{i+1}$ of Γ, respectively. If an edge e of Γ, belongs to neither $\mathcal{K}_i$ nor $\mathcal{K}_{i+1}$, then $\phi(e) \in \mathcal{A}_i$. In particular, Γ has no H-cells.*

Proof. The first assertion follows from Lemma 3.4, since the labels of the top and bottom of an i-sector are of the form $k_i \dots k_{i+1}$. Then, by the same lemma, the second assertion is true for the edges of all θ-cells since every maximal θ-band must connect $\mathcal{K}_i$ and $\mathcal{K}_{i+1}$. Since $i \neq 1$ and the labels of the boundary edges of H-cells belong to $\mathcal{A}_1$, the H-cells of Γ have no edges in common either with the θ-cells of Γ or with the boundary $\partial\Gamma$. Now the minimality of the diagram Γ implies that Γ has no H-cells at all, and so the lemma is proved. □

The following lemma claims that every i-sector, $i \neq 1$, simulates the work of $\mathcal{S}(L) \cup \hat{\mathcal{S}}(L)$. It summarizes the assertions of Lemmas 6.1, 6.3, 6.9, and 6.16 from [23]. For the formulation (1) below, it is important that $\mathcal{S}$ (and $\mathcal{S} \cup \hat{\mathcal{S}}$) is an S-machine. The analog of this statement is false for Turing machines. (See [21] for a discussion.)

Lemma 3.9. *(1) Let Δ be an i-sector for some $i \neq 1$ with history $\theta_1 \dots \theta_d$. Assume that Δ has consecutive maximal θ-bands $\mathcal{T}_1, \dots \mathcal{T}_d$, and $k_iW_jk_{i+1}$ and $k_iW_j'k_{i+1}$ are the bottom and the top labels of $\mathcal{T}_j$, ($j =$*

$1, \dots, d$). *Let* U_j *(resp.* V_j, $i = 1, \dots d$*) be the copies of* W_j *(resp.* W'_j*) in the alphabet of the machine* $\mathcal{S} \cup \hat{\mathcal{S}}$. *Then* U_j, V_j *are admissible words for* $\mathcal{S} \cup \hat{\mathcal{S}}$, *and*

$$V_1 \equiv U_1 \cdot \theta_1, U_2 \equiv V_1, \dots, U_d \equiv V_{d-1}, V_d \equiv U_d \cdot \theta_d$$

(2) For every reduced computation $U \cdot h \equiv V$ *of* $\mathcal{S}$ *(of* $\hat{\mathcal{S}}$*) with* $|h| \geq 1$ *and for every* $i \in [1, N]$ *(for every* $i \in [2, N]$*), there exists an* i*-sector* Δ *with history* h *and without* H*-cells, whose bottom and top labels are* $k_i U' k_{i+1}$ *and* $k_i V' k_{i+1}$, *where* U' *(resp.* V'*) is the copy or the mirror copy of the word* U *(of* V*) in the alphabet* $\mathcal{A}_i$. *These copies are mirror copies iff* i *is even.*

We call an i-sector *accepted* if its top label $k_i V' k_{i+1}$ is the i-sector subword of the word Σ_0. For $i \neq 1$, the computation $V_1 \to \dots \to V_d$ of the machine $\mathcal{S} \cup \hat{\mathcal{S}}$ provided by Lemma 3.9 (1) for an accepted i-sector is accepting.

3.4 Replicas

Let Δ be an i-sector, where $i \neq 1$. Then by Lemma 3.8, for every $i' \neq 1$, one can relabel the edges of Δ (or of the mirror copy of Δ if $i - i'$ is odd) and obtain an i'-sector Δ', which is just a copy (or a mirror copy) of Δ. This symmetry and mirror symmetry are necessary for the diagram surgery we will utilize in the proof of Lemma 4.9 of Section 4. But one cannot construct such a copy of the i-sector if $i' = 1$ since there are no commutativity relations $\hat{\theta}_j \hat{a} = \hat{a} \hat{\theta}_j$ if $\hat{\theta}_j \in \mathcal{A}_1$ (see (3.5)). However one can construct an ersatz-copy of Δ called a replica, if Δ is an accepted sector. This construction will be used in Sections 4 and 5.

For $i' = 1$, we construct the *replica* of Δ as follows. (We assume below that i is odd, otherwise one first replaces Δ by its mirror copy.)

At first, the relations (3.4) and (3.5) make it possible to replace the maximal k_i-band $\mathcal{K}_i$ and k_{i+1}-band $\mathcal{K}_{i+1}$ of Δ by their copies $\mathcal{K}_1$ and $\mathcal{K}_2$, respectively. Similarly, we replace every maximal θ-band of Δ by its copy if θ is a command of the machine $\mathcal{S}$. Now let $\mathcal{T}$ be a maximal θ-band of Δ and θ a command of the machine $\hat{\mathcal{S}}$. This $\mathcal{T}$ consists of (θ, q)- and (θ, a)-cells. To construct the replica $\mathcal{T}'$ of $\mathcal{T}$ we take the 'copies' of (θ, q)-cells only (but no a-edges in these 'copies', the a-edges are contracted to vertices) and build $\mathcal{T}'$ from them by identifying θ-edges of neighboring cells in the order in which the original (θ, q)-cells appear in $\mathcal{T}$.


It remains to close up the holes between θ-bands ${\mathcal{T}'}_s$ and ${\mathcal{T}'}_{s+1}$ for consecutive $\mathcal{T}_s$ and $\mathcal{T}_{s+1}$. If the corresponding letters θ_s and θ_{s+1} are both commands of $\mathcal{S}$, then we just identify the top of the copy ${\mathcal{T}'}_s$ and the bottom of the copy ${\mathcal{T}'}_{s+1}$. This is possible since $\phi(\mathbf{top}\mathcal{T}_s) \equiv \phi(\mathbf{bot}\mathcal{T}_{s+1})$ by Lemma 3.9 (1). A similar identification works if θ_s and θ_{s+1} are both the commands of $\hat{\mathcal{S}}$.

Assume now that θ_s is a command of $\mathcal{S}$ and θ_{s+1} is a command of $\hat{\mathcal{S}}$ (or vice versa). Then $\phi(\mathbf{top}(\mathcal{T}_s)) \equiv \phi(\mathbf{bot}\mathcal{T}_{s+1})) \equiv k_i W_s k_{i+1}$, and the copy W'_s of W_s in the alphabet $\mathcal{A}_1$ is an admissible word by both machines $\mathcal{S}$ and $\hat{\mathcal{S}}$ since both commands θ_s^{-1} and θ_{s+1} are applicable to it. But the only common tape letters of these machines are the letters of the input alphabet of $\mathcal{S}$, and the only common state letters of these machines are the letters from the start vector $\vec{s}_1$ and the accept vector $\vec{s}_0$ of $\mathcal{S}$. Then by property (5) of Lemma 2.10, either θ_s^{-1} is the start rule or θ_s is the accept rule of $\mathcal{S}$.

In the latter case, we have $\phi(\mathbf{top}({\mathcal{T}'}_s)) \equiv \phi(\mathbf{bot}{\mathcal{T}'}_{s+1})) \equiv k_i W_s k_{i+1}$ since the accept words have no a-letters at all, and so identification of the top of the copy ${\mathcal{T}'}_s$ and the bottom of the copy ${\mathcal{T}'}_{s+1}$ is possible again. Let us consider the former case. Then the rule θ_s^{-1} is of the form $q_1 \to q'_1, q_2 \stackrel{\ell}{\to} q'_2, \dots, q_K \stackrel{\ell}{\to} q'_K$, where $(q_1, \dots, q_K) \equiv \vec{s}_1$. Therefore W'_s can have tape letters only between q_1 and q_2, i.e., $W'_s \equiv q_1 u q_2 q_3 \dots q_K$, where u is a word in the input alphabet.

Since the sector Δ is accepted and the rules of S-machines are invertible, we have that W'_s is accepted by the machine $\mathcal{S}(L) \cup \hat{\mathcal{S}}(L)$, and so it is accepted by $\mathcal{S}$ by Lemma 3.1. Hence the word u belongs to the language recognized by the machine $\mathcal{S}$, and therefore u is a word in the generators of H, and $u = 1$ in H by the definition of that language.

Now, to close up the hole between the bands ${\mathcal{T}'}_s$ and ${\mathcal{T}'}_{s+1}$, it suffices to paste in an H-cell π labeled by the cyclically reduced form of the word u between them because $\phi(\mathbf{top}({\mathcal{T}'}_s)) \equiv k_1 q_1 u q_2 q_3 \dots q_K k_2$ and $\phi(\mathbf{bot}{\mathcal{T}'}_{s+1}) \equiv k_1 q_1 q_2 q_3 \dots q_K k_2$.



Replica

The replica Δ' of the accepted i-sector Δ is constructed. To summarize our effort: we cut off the (θ, a)-cell and contract up the a-edges of (θ, q)-cells for every command θ of the machine $\hat{\mathcal{S}}$ from Δ, replace the maximal k_i- and k_{i+1}-bands by their k_1- and k_2-copies, replace all the labels of the remaining edges by their copies from the alphabet $\mathcal{A}_1$, and then close up all the holes by pasting in several H-cells. The result is the replica Δ', canonically obtained above.

Remark 3.10. (1) The i-sector Δ is a union of $K+1$ *subsectors* $\Gamma_1, \dots, \Gamma_{K+1}$, where every Γ_j is a trapezia with a base of length 2 and with height equal to the height of Δ. The subsector Γ_j has a common maximal q-band $\mathcal{C} = \mathcal{C}_{j+1}$ with Γ_{j+1} for $j = 1, \dots, K$. If i is odd (even), then Γ_2 (resp., Γ_K) is the *input subsector*. When we construct a replica, H-cells appear only in the input subsector of the replica.

Every Γ_j has its own replica Γ'_j, which is a subsector of Δ'. Similarly, every q- or θ-edge of $\mathcal{C}$ has a replica in the replica $\mathcal{C}'$ of $\mathcal{C}$ in Δ'. Also every vertex o of $\mathcal{C}$ belongs to either a q-edge or a θ-edge since the boundary of a (θ, q)-cell has no two consecutive a-edges by Lemma 2.10(5); and so o has a replica o'.

(2) The replica of an i-sector is not necessarily a minimal diagram.

3.5 Discs

Given a diagram Δ, one can construct a planar graph whose vertices are the hubs of this diagram plus one improper vertex outside Δ, and the edges are maximal k-bands of Δ. Lemma 3.11 says that if the number L of sectors of a hub is large enough, then the degree of the improper vertex is positive provided the graph has at least one proper vertex; and therefore the homomorphism $H \to G$ turns out to be injective.

A *disc diagram (or a disc)* is a (sub)diagram Δ such that (1) it has exactly one hub Π (2) there are no θ-edges on the boundary $\partial\Delta$ (3) there are no H-cells of Δ having an edge on $\partial\Delta$. In particular, a hub is a disc diagram.


Let us consider a disc diagram Δ with hub Π. Denote by $\mathcal{K}_1, \dots, \mathcal{K}_L$ the $k_1, \dots, k_L$-bands starting on the hub Π. Since the hub relation has only one letter k_i for every i, these k-bands have to end on $\partial\Delta$. It therefore follows from Lemma 3.4 that for every i, the bands $\mathcal{K}_i$ and $\mathcal{K}_{i+1}$ bound, together with $\partial\Delta$ and $\partial\Pi$, either a subdiagram Ψ_i having no cells corresponding to any non-trivial relation of G (in this case the bands $\mathcal{K}_i$ and $\mathcal{K}_{i+1}$ are also trivial), or Ψ_i is a trapezium, and the boundary label of Δ has exactly one letter from $\mathcal{Q}_i$ for every $i \in [1, \dots, N]$.

Similarly, consider two hubs Π_1 and Π_2 in a minimal diagram, connected by a k_i-band $\mathcal{K}_i$ and a k_{i+1}-band $\mathcal{K}_{i+1}$, where $(i, i+1) \neq (1, 2)$, and there are no other hubs between these k-bands. These bands, together with $\partial\Pi_1$ and $\partial\Pi_2$, bound either a subdiagram Ψ_i having no cells, or a trapezium Ψ_i. The former case is impossible since in this case the hubs have a common k_i-edge and they are mirror copies of each other contrary to the reducibility of minimal diagrams. We want to show that the latter case is not possible either.

Indeed, in the latter case Ψ_i is an accepted trapezium since the i-sector subword of Σ_0 is $k_i w k_{i+1}$, where w is a (mirror) copy of the accept word of the machine $\mathcal{S}$. Therefore, according to Subsection

3.4, a replica Ψ_1 of Ψ_i (as well as the (mirror) copies Ψ_j for every $j \in [2, \dots, N]$) can be constructed. Then one can construct a spherical diagram Γ from Π_1, Π_2, and the diagrams Ψ_j $(j = 1, \dots, N)$. There are two subdiagrams Γ' and Γ'' of Γ with common boundary: Γ', being a copy of a subdiagram of the original diagram, is made of Π_1, Π_2, and $\Psi = \Psi_i$, and Γ'' is a union of all Ψ_j-s with $j \neq i$. Hence the subdiagram Γ' of the original diagram can be replaced by diagram of lower type with the same boundary label because Γ'' has no hubs. This contradicts the minimality of the original diagram.

Thus, any two hubs of a minimal diagram are connected by at most two k-bands, such that the subdiagram bounded by them contains no other hubs. This property makes the hub graph of a minimal diagram (where maximal k-bands play the role of edges connecting hubs) hyperbolic (in a sense) since the degree L of every vertex (=hub) is high (≥ 40). Below we give a more precise formulation (proved for diagrams with such a hub graph, in particular, in [20], Lemma 3.2).



Lemma 3.11. *If a minimal diagram over the group G contains a least one hub, then there is a hub Π in Δ such that $L-3$ consecutive maximal k-bands $\mathcal{B}_1, \dots \mathcal{B}_{L-3}$ start on Π, end on the boundary $\partial\Delta$, and for any $i \in [1, L-4]$, there are no discs in the subdiagram Γ_i bounded by $\mathcal{B}_i$, $\mathcal{B}_{i+1}$, $\partial\Pi$, and $\partial\Delta$.*

Corollary 3.1. *The canonical homomorphism $H \to G$ given by Lemma 3.2 is injective.*

Proof. Assume that a word w in the generators $a_1, ..., a_m$ of the group H is equal to 1 in G. Then by van Kampen's Lemma, there is a minimal diagram Δ over G whose boundary label is w. Since w has no q-letters, Δ has no hubs by Lemma 3.11. By Lemma 3.4, Δ contains neither q- nor θ-annuli, and so it has neither (θ, q)-cells nor (θ, a)-cells, because w has neither q- nor θ-letters. Hence this diagram can contain H-cells only. Since the boundary labels of H-cells are trivial in H, the boundary label w is trivial in H too by van Kampen's Lemma, and so the homomorphism is injective. □

.

4 Comparison of paths in diagrams

The main lemma of this section is Lemma 4.9 which says that given a diagram Δ, one can cut off a subdiagram having one hub (and cells of lower rank) such that the perimeter of the remaining diagram does not exceed $|\partial\Delta|$. This makes it possible to use induction on the number of hubs in the next section. To carry out the surgery of Lemma 4.9 we must be able to compare the lengths of paths in various particular diagrams and to 'finish' some 'unfinished' diagrams.

4.1 Paths in sectors

We will consider the words in the generators of the group G and will modify the length function on this set. This modification is helpful in subsequent subsections, in particular, we cannot prove an analog of Lemma 4.6 using the standard length $||\ ||$.

The standard length $||\ ||$ of a word (of a path) will be called its *combinatorial length*. From now on we use the word *length* for the modified length. We set the length of every q-letter equal 1, and the length of every a-letter equal to a small enough number $\delta > 0$ so that

$$\delta < (3N)^{-1}. \tag{4.7}$$

If a word v has s θ-letters, t a-letters, and no q-letters, then

$$|v| = s + \delta \max(0, t - s)$$

by definition. For example, the word read between two q-letters of a (q, θ)-relation has length 1 since it has one θ-letter and at most one a-letter by formulas (3.4, 3.5) and property (3) of Lemma 2.10.


An arbitrary word w is a product $v_0 q_1 v_1 q_2 \dots q_m v_m$, where $q_1, \dots, q_m$ are q-letters and the words $v_0, \dots v_m$ have no q-letters. Then, by definition, $|w| = m + \sum_{j=0}^{m} |v_j|$. *The length of a path* in a diagram is the length of its label. The *perimeter* $|\partial\Delta|$ of a van Kampen diagram is similarly defined by a shortest cyclic decomposition of the boundary $\partial\Delta$. It follows from this definition that for any product $s = s_1 s_2$ of two words or paths, we have $|s| \le |s_1| + |s_2|$, and $|s| = |s_1| + |s_2|$ if s_2 starts or s_1 ends with a q-letter.

If a path p starts at a vertex o and ends at o', we will write $o = p_-$ and $o' = p_+$.

Lemma 4.1. *Let Δ be a trapezium bounded by two maximal q-bands $\mathcal{C}$ and $\mathcal{C}'$ and having no H-cells. Assume that $\mathcal{C}$ has no a-edges. Let o be vertex lying on both $\mathcal{C}$ and the top of the trapezium Δ, and let o' belong to $\mathcal{C}'$ and to the bottom of Δ. Assume that a path t connects o and o' and has no q-edges. Then the vertices o and o' can be connected in Δ by a path t' such that $|t'| \le |t|$ and $t' = t_1 t_2 t_3$ where t_1 and t_3 are parts of the sides of $\mathcal{C}$ and $\mathcal{C}'$, respectively, and t_2 consists only of a-edges.*

Proof. Since the path t has no q-edges, it follows from the assumption of the lemma that every maximal θ-band $\mathcal{T}$ of Δ has exactly two (θ, q)-cells (the first one and the last one). Therefore $\mathcal{C}$ and $\mathcal{C}'$ can be connected along $\mathcal{T}$ by a path x consisting of a-edges only.

We denote by t_2 a shortest path among such x-s. Then we define t_1 (t_3) as the shortest subpath of the side of $\mathcal{C}$ (of $\mathcal{C}'$) connecting o and $(t_2)_-$ ($(t_2)_+$ and o').



Assume that there is an a-band $\mathcal{A}$ starting with an a-edge of t_2 and ending with an a-edge e of $\partial\mathcal{C}'$. Then e belongs to some path ye where y consists of a-edges and connects $\mathcal{C}$ and $\mathcal{C}'$. Notice that every maximal a-band crossing the path y must cross t_2 because it cannot cross $\mathcal{A}$, and $\mathcal{C}$ has no a-edges. Hence $|y|_a \le |t_2|_a - 1$, contrary to the minimality in the choice of t_2.

Thus every maximal a-band $\mathcal{A}$ crossing t_2 must connect the top and the bottom of the trapezium Δ, and therefore the path t must cross every such a-band $\mathcal{A}$. Also t must cross every maximal a-band

starting on t_3 whence $|t|_a \geq |t_2|_a + |t_3|_a = |t'|_a$. Since the path $|t|$ must cross every maximal θ-band of Δ we also have the inequality $|t|_\theta \geq |t_1|_\theta + |t_3|_\theta = |t'|_\theta$. Now it follows from the definition of path length that $|t'| \leq |t|$ as required. □


Lemma 4.2. *Let Δ be an accepted i-sector with index $i \neq 1$ bounded by a maximal k_i-band $\mathcal{C}$ and a k_{i+1}-band $\mathcal{C}'$. Let o_1 and o_2 be two vertices lying on $\mathcal{C}$ and $\mathcal{C}'$, respectively. Assume that o_1 and o_2 are connected by a path t, in Δ. Then the replicas o_1' and o_2' of the vertices o_1 and o_2 in the replica Δ' of Δ can be connected by a path t' such that $|t'| \leq |t|$.*

Proof. First of all, one may assume that no maximal q-band $\mathcal{C} = \mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_{k+2} = \mathcal{C}'$ is crossed by the path t twice. Indeed, otherwise t has a subpath s of the form ezf, where e and f are q-edges of some $\mathcal{C}_j$ separated in this band by m (θ, q)-cells for some $m \geq 0$. Therefore the path z must cross at least m maximal θ-bands whence $|ezf| \geq m + 2$. But the vertices e_- and f_+ can be connected along $\mathcal{C}_j$ by a path of length m (see the example after the definition of length $|*|$), and so the path t can be shortened.

Thus the path t is a product $t = t_1 \dots t_{k+1}$, where each t_j connects a vertex $o(j)$ lying on $\mathcal{C}_j$ with a vertex $o(j+1)$ lying on $\mathcal{C}_{j+1}$, and for every $j = 1, \dots, k$, either t_{j+1} starts or t_j ends with a q-edge, and so $|t| = \sum_{j=1}^{k+1} |t_j|$. As in the previous paragraph, we have that each of the t_j-s crosses every θ-band at most once. (Consider ezf, where e and f are θ-edges of the same θ-band.) Now using the notation of Remark 3.10, it suffices to consider the replica Γ_j' of the subsector Γ_j and to find a path t_j' connecting the replicas $o'(j)$ and $o'(j+1)$, with $|t_j'| \leq |t_j|$.

We may assume that i is odd. (If i is even one should use a mirror argument.)

We first consider the path t_2 crossing the input subsector Γ_2, assuming that t_2 has no q-edges, since the q-edges (if any) can be attributed to the subpaths t_1 and t_3. By property (6) of Lemma 2.10, $\mathcal{C}_2$ has no a-edges. Hence, by Lemma 4.1 applied to a subtrapezium of Γ_2 containing t_2, we may assume that $t_2 = s_1 s_2 s_3$, where s_1 and s_3 are the subpaths of the top or bottom paths of q-bands $\mathcal{C}_2$ and $\mathcal{C}_3$, respectively, and s_2 goes along the top or bottom of a maximal θ-band $\mathcal{T}$. For both paths s_1 and s_3, we have paths s_1' and s_3' of the same length lying on the boundaries of the q-bands $\mathcal{C}_2'$ and $\mathcal{C}_3'$ of the replica Γ_2' and connecting the replicas of the vertices $(s_1)_\pm$ and $(s_3)_\pm$, respectively. The vertices $(s_1')_+$ and $(s_3')_-$ are either connected by a copy s_2' of s_2 (if the θ-band $\mathcal{T}$ was copied when we constructed the replica Δ') or $(s_1')_+ = (s_3')_-$ (if the corresponding θ-band of Δ' has no a-edges). It follows that in any case we have $|t_2'| \leq |t_2|$ for $t_2' = s_1' s_2' s_3'$.



Assume now that $j \neq 2$. The subsector Γ_j has no H-cells by Lemma 3.8, and so it is a union of alternating subtrapezia $\Gamma_{j1}, \Gamma_{j2}, \dots$ whose histories are words either in the alphabet Θ or in $\hat{\Theta}$. Let $t_j = x_1 \dots x_d$, where every x_k belongs to Γ_{jk}. If the history of the trapezia Γ_{jk} is a word over Θ, then we have the copy Γ_{jk}' of Γ_{jk} in Δ', and a subpath $x = x_k$ of t_j lying in Γ_{jk} has a copy x' in Γ_{jk}'. If the history is a word over $\hat{\Theta}$, then Γ_{jk}' has no a-edges, and for every subpath x of t_j lying in Γ_{jk}, we can construct a corresponding subpath x' in Γ_{jk}' which copies only q- and θ-edges of x, but ignores the


a-edges of x. Since there are no a-edges in the common boundaries of the neighbors Γ'_{jk} and $\Gamma'_{j,k+1}$, we have $(x'_k)_+ = (x'_{k+1})_-$ for every $k = 1, \dots, d-1$, and we obtain $|t'_2| \le |t_2|$ for the path $t'_2 = x'_1 \dots x'_d$.

Now the required path t' is obtained, and the lemma is proved. □

Assume that we have a minimal diagram Γ with a cyclically reduced boundary label over the group $G(\mathcal{S} \cup \hat{\mathcal{S}}, L)$, which is separated by a maximal k_i-band (or k_{i+1}-band) $\mathcal{C}$ in two parts $\tilde{\Delta}$ and Δ such that Δ is an accepted i-sector. Assume that every maximal θ-band $\mathcal{T}_1, \dots, \mathcal{T}_m$ of Γ crosses Δ, the bottom x of $\mathcal{T}_1$ is a part of the boundary $\partial\Gamma$, and $\phi(x)$ is a suffix of the subword $k_{i-1} \dots k_i \dots k_{i+1}$(a prefix of the subword $k_i \dots k_{i+1} \dots k_{i+2}$, respectively) of the word Σ_0. Also we assume that for every $j \in [2, m]$, the trimmed bottom $\mathbf{tbot}\mathcal{T}_j$ is a subpath of the top $\mathbf{top}\mathcal{T}_{j-1}$. Below we call such a diagram *unfinished* if $i, i-1 \ne 1$ (if $i, i+1 \ne 1$, respectively).

Lemma 4.3. *The part $\tilde{\Delta}$ of the unfinished diagram Γ can be embedded in an $i-1$-sector (respectively, $i+1$-sector) ∇ which is a mirror copy of the i-sector Δ.*

Proof. Without loss of generality we assume that $\mathcal{C}$ is a k_i-band in the definition of unfinished diagram Γ. We construct ∇ by induction on the length m of the history $h \equiv \theta_1 \dots \theta_m$ of the i-sector Δ.



Let Π be the intersection cell for $\mathcal{T}_1$ and $\mathcal{C}$. Then $\mathcal{T}_1$ consists of Π and two subbands $\mathcal{T}_1'$ and $\mathcal{T}_1''$, where $\mathcal{T}_1'$ belongs to $\tilde{\Delta}$ and $\mathcal{T}_1''$ belongs to Δ. Let $\Pi(1)'$ and $\Pi(1)''$ be the neighboring cells for Π in $\mathcal{T}_1'$ and in $\mathcal{T}_1''$, respectively. Then $\Pi(1)'$ (if it exists) is a mirror copy of $\Pi(1)''$ with boundary label in the alphabet $\mathcal{A}_{i-1}$. Indeed, these two cells are determined by the same history θ_1 and mirror q-letters of the word Σ_0 as this follows from Relations (3.4, 3.5). A similar argument shows that if $\Pi'(1)$ has a neighbor $\Pi'(2)$ (where $\Pi'(2) \ne \Pi$) in $\mathcal{T}_1'$, then the cell $\Pi''(1)$ has a neighbor $\Pi''(2)$ (where $\Pi''(2) \ne \Pi$) in $\mathcal{T}_1''$, and $\Pi'(2)$ is a mirror copy of $\Pi''(2)$ with boundary label in the alphabet $\mathcal{A}_{i-1}$. By induction, we obtain that $\mathcal{T}_1'$ is a mirror copy of a subband of $\mathcal{T}_1''$ starting with Π. Hence one can extend $\mathcal{T}_1'$ (and the subdiagram $\tilde{\Delta}$) and obtain a θ-band $\mathcal{T}_1^{\nabla}$ which is a mirror copy of $\mathcal{T}_1''$.

Since the trimmed bottom of the band $\mathcal{T}_2$ is a subpath of the top of $\mathcal{T}_1$, one can similarly subdivide $\mathcal{T}_2$ in $\Pi(2)$, $\mathcal{T}_2'$, $\mathcal{T}_2''$, and prove that $\mathcal{T}_2'$ is the mirror copy of a subband of $\mathcal{T}_2''$, starting with the cell $\Pi(2)$ and having boundary label over the alphabet $\mathcal{A}_{i-1}$. ($\mathcal{T}'_2$ and $\mathcal{T}''_2$ may include (θ, q)-cells and also (θ, a)-cells.) Therefore there is an extension $\mathcal{T}_2^{\nabla}$ of $\mathcal{T}_2'$, and this extension is a mirror copy of $\mathcal{T}_2''$. Then by induction, we construct θ-bands $\mathcal{T}_3^{\nabla}, \dots, \mathcal{T}_m^{\nabla}$, and these θ-bands together with the q-band $\mathcal{C}$ form the required $i-1$-sector ∇. □

Lemma 4.4. *Let Δ be a minimal diagram with boundary path $x_1 y_1 x_2 y_2$, where*

(1) $\phi(y_1)$ is a subword $k_j \dots k_i$ of Σ_0, and this subword does not contain the letter k_1;

(2) x_1 and x_2 are sides of the maximal k_j-band $\mathcal{K}_j$ and k_i-band $\mathcal{K}_i$ starting on y_1;

(3) every cell of Δ belongs to one of the maximal θ-bands $\mathcal{T}_1, \dots, \mathcal{T}_m$ of Δ;

(4) (a) either each of the bands $\mathcal{T}_1, \dots, \mathcal{T}_m$ crosses $\mathcal{K}_j$ or (b) each of them crosses $\mathcal{K}_i$;

(5) the trimmed bottom path of $\mathcal{T}_1$ is a subpath of $y_1^{\pm 1}$, and the trimmed bottom path of $\mathcal{T}_l$ is a subpath of the top path of $\mathcal{T}_{l-1}$ for every $l = 2, \dots, m$;


(6) one can construct a diagram $\bar{\Delta}$ with boundary $\bar{x}_1\bar{y}_1\bar{x}_2\bar{y}_2$, and $\bar{\Delta}$ satisfies the analogs of properties (1)-(5), but $\phi(\bar{y}_1) \equiv k_{j-1} \dots k_i$ in case 4(a) ($\phi(\bar{y}_1) \equiv k_j \dots k_{i+1}$ in case 4(b)), and Δ is embeddable in $\bar{\Delta}$ so that the k_j-band $\mathcal{K}_j$ and the k_i-band $\mathcal{K}_i$ remain maximal in $\bar{\Delta}$.

Then in case (4)(a) (in case 4(b)), there exists a diagram Δ' over the group G_1 with boundary path $x'_1y'_1x'_2y'_2$ such that $\phi((y'_1)^{-1})$ is the subword $k_{2j-i} \dots k_j$ (respectively, $k_i \dots k_{2i-j}$) of Σ_0, x'_1 and x'_2 are sides of the maximal k_j-band $\mathcal{K}_j$ and k_{2j-i}-band $\mathcal{K}_{2j-i}$ (of the maximal k_i-band $\mathcal{K}_i$ and k_{2i-j}-band $\mathcal{K}_{2i-j}$ starting on y'_1), the labels of x'_1 and x'_2 are copies of $\phi(x_1)$ and $\phi(x_2)$, resp., and $|y'_2| \le |y_2|$.

(The subscripts of k-bands are taken modulo L.)

Proof. We consider the case (4)(b) only. The maximal k-bands $\mathcal{K}_j, \mathcal{K}_{j+1} \dots \mathcal{K}_{i+1}$ subdivide the diagram $\bar{\Delta}$ in the subdiagrams Γ_l-s, where Γ_l is bounded by $\mathcal{K}_l$ and $\mathcal{K}_{l+1}$ and Γ_l includes these k-bands ($l = j, \dots, i$).

Every maximal θ-band of Δ having a cell in Γ_{i-1}, must cross both bands $\mathcal{K}_i$ and $\mathcal{K}_{i+1}$ of $\bar{\Delta}$. Therefore the parts of these bands in Γ_{i-1} and Γ_i form an unfinished diagram whose i-sector is a subdiagram Δ_i of Γ_i. By Lemma 4.3, the subdiagram Γ_{i-1} is embedded into the mirror copy Δ_{i-1} of Δ_i. Similarly, Γ_{i-2} is embeddable into a mirror copy of Δ_{i-1} which is the copy of Δ_i (we denote this copy by Δ_{i-2}) ,..., Γ_j is embeddable into the copy (or mirror copy) Δ_j of Δ_i.



Let $y_2 = t_{i-1} \dots t_j$, where t_l passes through Γ_l (and Δ_l) for $l = i-1, \dots, j$. Since every subpath t_l connects two vertices on the k-bands of the l-sector Δ_l, we can construct the mirror copy Δ'_{2i-l-1} (which is a $2i-l+1$-sector) of Δ_l or the replica of Δ_l (if $2i-l-1=1$), and the copies of the vertices $(t_l)_\pm$ are connected in Δ'_{2i-l-1} by a path t'_{2i-l-1} with $|t'_{2i-l-1}| \le |t_l|$ by Lemma 4.2. The desired diagram Δ' embeds in the union of these Δ'_{2i-l-1}-s, and $y'_2 = t'_i \dots t'_{2i-j-1}$. □

4.2 Shortcuts

In this subsection we show that Lemma 4.4 helps cut off a hub from a diagram using a 'shortcut'. But at first we consider a few statements about simpler pieces (without hubs) of diagrams over the group G.

Lemma 4.5. *Suppose a diagram Δ over G has a q-band $\mathcal{C}$ starting and ending on $\partial\Delta$, and p is a side of $\mathcal{C}$. Assume that no θ-band crosses $\mathcal{C}$ twice in Δ and that there is a factorization xy of the boundary path of Δ such that $x_- = p_-, x_+ = p_+$. Then $|p| \le |x|$, $|\partial\mathcal{C}| \le |\partial\Delta|$, and $|xp^{-1}| \le |\partial\Delta|$.*

Proof. On the one hand, the length $|p|$ of p is equal to the number m of θ-cells in $\mathcal{C}$, since every cell of $\mathcal{C}$ has one θ-edge and at most one a-edge on p by condition 3 of Lemma 2.10 and the definition of (θ, q)-relations. On the other hand, every maximal θ-band crossing $\mathcal{C}$ must terminate on x, since it does not cross $\mathcal{C}$ twice. It follows that $|x| \ge m$, and so $|p| \le |x|$. Similarly, we obtain inequalities $|\partial\mathcal{C}| \le |\partial\Delta|$,


and $|xp^{-1}| \le |\partial\Delta|$. (We take into account the definition of length and the fact that the sides of the band $\mathcal{C}$ are separated by two q-edges lying on $\partial\Delta$.) □

A maximal θ-band of a diagram Δ called a *rim band* if its start and end θ-edges as well as its top or its bottom lies on the boundary path of Δ.

Lemma 4.6. *Suppose Δ is a diagram over G with a rim band $\mathcal{T}$ having at most N (θ, q)-cells. Denote by Δ' the subdiagram $\Delta \backslash \mathcal{T}$. Then $|\partial\Delta| - |\partial\Delta'| \ge 1/2$.*

Proof. Let s be the top side of $\mathcal{T}$ and $s \subset \partial\Delta$. Note that by our assumptions the difference between the number of a-edges in the bottom s' of $\mathcal{T}$ and the number of a-edges in s cannot be greater than $2N$ since every (θ, q)-cell has at most two a-edges. However, Δ' is obtained by cutting off $\mathcal{T}$ along s', and its boundary contains two fewer θ-edges than Δ. Thus one can compare the boundaries of Δ and Δ' as follows. There is a one-to-one correspondence between the q-edges of these boundaries, and to extend this correspondence to the θ- and a-edges of the boundaries, one should remove two θ-edges from $\partial\Delta$ and add at most $2N$ a-edges. Therefore it follows from the definition of length and by Inequality (4.7), that $|\partial\Delta| - |\partial\Delta'| \ge 2(1-\delta) - 2N\delta > 1/2$. □

Lemma 4.7. *Suppose a diagram Δ has two cells: an H-cell Π and a (θ, a)-cell π which have a common a-edge e, ep is the boundary of Π and $efe'f'$ is the boundary of π, where $\phi(e) \equiv \phi(e')^{-1} \equiv a$ and $\phi(f) \equiv \phi(f')^{-1} \equiv \theta$ for a θ-letter θ. Then there is a diagram Δ' with the same boundary label as Δ composed of Π and a θ-band $\mathcal{T}$, and p is the side of $\mathcal{T}$.*



Proof. The letter a commutes with θ, and therefore every letter of the boundary label of Π commutes with θ. Hence one can construct a θ-band $\mathcal{T}$ with boundary $g'p^{-1}gp'$, where $\phi(p') \equiv \phi(p)$, and $\phi(g) \equiv \phi(g')^{-1} \equiv \theta^{-1}$. If we attach the band $\mathcal{T}$ to Π along the path p and remove π, we obtain the required diagram Δ'. □

Let Π be a hub of a minimal diagram Δ given by Lemma 3.11. Using the notation of that lemma, we recall that the subdiagrams Γ_i and Γ_{i+1} intersect along the k-band $\mathcal{B}_{i+1}$ $(i = 1, \dots, L-5)$. We denote by Ψ the minimal subdiagram containing all the Γ_i-s for $i = 1, \dots, L-4$. The boundary path of Ψ is $x'x''$, where x' is composed of the sides of $\mathcal{B}_1$, $\mathcal{B}_{L-3}$, and a subpath of $\partial\Pi$, while x'' is a subpath of $\partial\Delta$.

Lemma 4.8. *One can construct a minimal diagram Ψ' over G_1 with boundary path $x'\bar{x}$ such that (1) Ψ' includes the bands $\mathcal{B}_1$ and $\mathcal{B}_{L-3}$ (2) $|\bar{x}| \le |x''|$, (3) the subdiagram Ψ' has no maximal q-bands except for the q-bands $\mathcal{B}'_i$ $(i = 1, \dots, L-3)$ starting on x', (4) every maximal θ-band of Ψ' crosses either the band $\mathcal{B}'_1 = \mathcal{B}_1$ or the band $\mathcal{B}'_{L-3} = \mathcal{B}_{L-3}$, (5) the subdiagram Ψ' has no H-cells between the pair of k-bands $\mathcal{B}'_i$ and $\mathcal{B}'_{i+1}$, unless this pair is a pair of k_1- and k_2-bands.*

Proof. Let Ψ' be a minimal diagram with boundary of the form $x'\bar{x}$ which satisfies conditions (1) and (2) and has minimal $|\bar{x}|$. Since Ψ satisfies conditions (1) and (2) with $\bar{x} = x'$, such Ψ' exists. Clearly the path $\bar{x}$ has no loops. If the diagram Ψ' has a maximal q-band $\mathcal{C}$ which does not start/terminate on Π, then one can cut off $\mathcal{C}$ and shorten $\bar{x}$ by Lemma 4.5. So Ψ' satisfies condition (3).

Assume that the diagram Ψ' does not satisfy condition (4) of the lemma. Then by Lemma 3.4, we have a θ-band of Ψ' starting and terminating on $\bar{x}$. It follows that there is a θ-band $\mathcal{T}$ starting and terminating on $\bar{x}$, such that the subdiagram Φ bounded by $\mathcal{T}$ and a part y of $\bar{x}$ has no non-trivial θ-bands, i.e., it contains only H-cells. Two H-cells of Φ cannot have a common edge since otherwise they could be replaced by one H-cell contrary to the minimality of Ψ'. It follows that every H-cell π of Φ has a common edge with $\mathcal{T}$ because the path $\bar{x}$ has no loops.

Thus we have a series of H-cells $\pi_1, \dots, \pi_s$ in Φ with boundaries $y_i z_i$ $(i = 1, \dots, s)$, where y_i is a part of y (or y_i is empty) and z_i belongs to the side z of $\mathcal{T}$. If $\sum |\phi(z_i)|_a \le 2$, then $\sum |z_i| \le 2\delta$ since every z_i is a product of a-edges. Therefore $|z| \le |\bar{x}| + 2\delta$. It follows from Lemma 4.6 that if we remove all π_i-s and then cut off the band $\mathcal{T}$, then we decrease the length of $\bar{x}$ since $2\delta < 1/2$; a contradiction.

Hence $\sum |\phi(z_i)|_a \ge 3$, and so at least one of π_i-s has a common edge with a (θ, a)-cell of $\mathcal{T}$. (We recall that $\mathcal{T}$ intersects each of the maximal q-bands of Ψ' starting on x'' at most once by Lemma 3.4, and so at most two of the (θ, q)-cells of $\mathcal{T}$ have a-edges with $a \in \mathcal{A}_1$, and each of these two (θ, q)-cells can have at most one a-edge with label from $\mathcal{A}_1$.) Therefore we can apply Lemma 4.7 to replace the (θ, a)-cell by a θ-band passing around the cell π_i. This modification of the band $\mathcal{T}$ decreases the number of H-cells in Φ, since one of the H-cells gets moved over the band $\mathcal{T}$.

The modified diagram can be non-minimal, but our surgery preserves q-bands and keeps the property that every q-band and θ-band have at most one common (θ, q)-cell. So sooner or later, this trick makes the inequality $\sum |\phi(z_i)|_a \le 2$ true, and one can decrease $\bar{x}$, as was explained above. If one replaces the obtained diagram by a minimal one, then condition (1) still holds since the maximal q-bands $\mathcal{B}_1$ and $\mathcal{B}_{L-3}$ are completely determined by the boundary as this follows from Lemma 3.4. This contradicts the assumption on the minimality of $|\bar{x}|$. Thus property (4) holds.

If Ψ' has H-cells in the subdiagram Γ'_i between the pair of k-bands $\mathcal{B}'_i$ and $\mathcal{B}'_{i+1}$ which are not a pair of k_1- and k_2-bands, then these H-cells (with labels over the alphabet $\mathcal{A}_1$) cannot have common edges with the maximal θ-bands of Γ'_i since the θ-bands of Γ'_i must intersect either $\mathcal{B}'_i$ or $\mathcal{B}'_{i+1}$ by (4). This implies that Γ'_i has no H-cells at all because the path $\bar{x}$ has no loops. The lemma is proved. □

Lemma 4.9. *If a minimal diagram Δ has a hub, then the (cyclic shift of the) boundary path of Δ can be factorized as pp' so that the subpath p starts and ends with q-edges and there is a simple path z in Δ with $z_- = p_-, z_+ = p_+$ such that the subdiagram bounded by the loop pz^{-1} has exactly one hub Π and the label $\phi(z)$ is equal in the group G_1 to a word of length $< |p|$.*

Proof. We may assume that a hub Π is chosen in Δ according to Lemma 3.11. Let $x'x''$ be the boundary path of Ψ as in Lemma 4.8. We will look for the path z in the minimal subdiagram Δ' obtained after removing Ψ and Π from Δ. Therefore to prove the lemma, one may assume using the notation of Lemma 4.8, that $\Psi' = \Psi$ and $\bar{x} = x''$, i.e., the subdiagram Ψ itself has properties (3), (4), and (5) from Lemma 4.8. (We do not know if the diagram $\Psi' \cup \Pi \cup \Delta'$ is still minimal but we will use the minimality of Ψ' only.)


There is $d \ge 0$ such that there exist exactly d maximal θ-bands of Ψ crossing each of the k-bands $\mathcal{B}_1, \dots, \mathcal{B}_{L-3}$. This implies that the initial subbands $\mathcal{B}_i[d]$ of length d in all $\mathcal{B}_i$-s $(i = 1, \dots, L-3)$ are copies of each other under the shifts of the indices in their boundary labels.

Let T be the set of remaining maximal θ-bands of Ψ, i.e., every band of T intersects exactly one of the bands $\mathcal{B}_1$, $\mathcal{B}_{L-3}$. It follows from Lemma 3.4 for Ψ that there is an integer l $(1 \le l < L-3)$ such that no θ-band of T crossing $\mathcal{B}_1$ crosses $\mathcal{B}_{l+1}$ and no θ-band of T crossing $\mathcal{B}_{L-3}$ crosses $\mathcal{B}_l$. We have either $(L-3) - l < (L-3)/2$ or $(l+1) - 1 < (L-3)/2$ since L is even. Without loss of generality we choose the former inequality, and so $l \ge (L-2)/2$.

If $\mathcal{B}_i$ is k_1-band for some $i \le 6$, then we will consider a smaller subdiagram bounded by $\mathcal{B}_7$ and $\mathcal{B}_{L-3}$ instead of Ψ. (Respectively, we change the complimentary minimal subdiagram Δ'.) If none of $\mathcal{B}_1, \dots, \mathcal{B}_6$ is a k_1-band, then we do not change Ψ. Thus, in any case we can reindex the k-bands and assume that Ψ is bounded by $\mathcal{B}_1$ and $\mathcal{B}_{L-r}$ for some $r \le 9$ and that the bands $\mathcal{B}_1, \dots, \mathcal{B}_{r+3}$ are not k_1-bands. Let them be k_i-,..., $k_{i\pm(r+2)}$-bands for some i. We will assume that $\mathcal{B}_{r+3}$ is a k_{i-r-2}-band.

Since $L \geq 40$, after such a reindexing $l \geq (L-2)/2 - 6 > 12 \geq r+3$, and so no θ-band from the set T crossing the band $\mathcal{B}_{L-3}$ crosses $\mathcal{B}_{r+3}$. We denote by Φ the part of the diagram Ψ bounded by $\mathcal{B}_{r+3}$ and $\mathcal{B}_1$. Let $\mathcal{T}_1^\Phi, \dots, \mathcal{T}_s^\Phi$ be the maximal θ-bands of Φ. Then by the choice of the subdiagrams Ψ and Φ, every cell of Φ belongs to one of these θ-bands, and each of these bands crosses the k_i-band $\mathcal{B}_1$. We will assume that $\mathcal{T}_1^\Phi$ is the closest band to the hub Π, and so on.

For every $j \geq 1$, at least one of the two q-edges of every (θ, q)-cell of $\mathcal{T}_i^\Phi$ belongs to the top of $\mathcal{T}_{j-1}^\Phi$ (to $\partial\Pi$ if $j = 1$) because Ψ has no maximal q-bands except for the bands starting on Π. Assume that a band $\mathcal{T}_j^\Phi$, starting with a (θ, q)-cell of $\mathcal{B}_1$, terminates with an (θ, a)-cell π having no a-edges on $\partial\mathcal{T}_{j-1}^\Phi$. Then an a-edge and one θ-edge of π lie on the boundary subpath x'' of Ψ, and so if one removes π from Ψ, the length of x'' does not increase because $\partial\pi$ has two θ-edges and two a-edges and all properties (3)-(5) hold for the remaining part of Ψ. Therefore we may assume that every edge f of $\mathbf{tbot}(\mathcal{T}_j^\Phi)$ belongs to $\mathbf{top}(\mathcal{T}_{j-1}^\Phi)$ (to $\partial\Pi$ for $j = 1$).



To construct the path z, we go along the side of the band $\mathcal{B}_2$ which is closer to $\mathcal{B}_1$, then go along the part of the boundary of the hub which is not part of $\partial\Psi$, and finally go to $\partial\Delta$ along the side of $\mathcal{B}_{L-r}$ which is closer to $\mathcal{B}_{L-r+1}$. Thus we have $z = z_1z_2z_3$ according to this definition, and respectively, $\phi(z) \equiv Z \equiv Z_1Z_2Z_3$. Let p be the subpath of $\partial\Delta$ and $\partial\Psi$ such that $p_- = z_-, p_+ = z_+$. Then the boundary path of Δ is of the form pp', for an appropriate p'.

We denote by Γ the part of the diagram Φ bounded by $\mathcal{B}_{r+3}$ and $\mathcal{B}_2$. Let y be the maximal common subpath of $\partial\Gamma$ and p with $y_- = p_- = z_-$ that does not cross the band $\mathcal{B}_{r+3}$. We may apply Lemma 4.4 to the pair (Γ, Φ) and obtain a new diagram Γ' over G_1 given by that lemma. One of the four boundary sections of Γ' is a subword $k_{i-1}...k_{i+r+1}$ of the word Σ_0.

Note that $\mathcal{B}_{L-r}$ is a k_{i+r+1}-band since we take the indices of the k-letters modulo L. By Lemma 4.4, Γ' has a loop with label of the form $Z_1Z_2Z'Y'$, where Z' is the copy of the word written along the band $\mathcal{B}_{r+3}$, and $|Y'| \leq |y|$. Hence $Z = (Y')^{-1}(Z')^{-1}Z_3$ in G_1. Therefore to prove that Z is equal in G_1 to a word of length $\leq |p|$, it suffices to prove that the word $(Z')^{-1}Z_3$ is equal in G_1 to a word of length $< |x|$, where $p = yx$.

We observe that the words Z' and Z_3 have equal prefixes of length d since $\mathcal{B}_{L-r}[d]$ is a copy of $\mathcal{B}_{r+3}[d]$. Therefore the word $(Z')^{-1}Z_3$ is equal to $(\bar{Z}')^{-1}\bar{Z}_3$, where $(\bar{Z}')^{-1}$ copies the label of the part of the side of $\mathcal{B}_{r+3}$ crossed by the θ-bands from the set T only, and $\bar{Z}_3$ is the label of the part of z_3 crossed by the θ-bands of T only. We denote the union of these two subsets of T by $\bar{T}$. The length of $(\bar{Z}')^{-1}\bar{Z}_3$ does not exceed the number $|\bar{T}|$ of bands in $\bar{T}$ since neither of the bands from T crosses both $\mathcal{B}_{r+3}$ and $\mathcal{B}_{L-r}$. By Lemma 3.4, every band of $\bar{T}$ must end on the subpath x. Hence $|(\bar{Z}')^{-1}\bar{Z}_3| \leq |x|$. In fact this inequality is strict since $L - r > r + 3$ (as $r \leq 9$) and so x must include some q-edges as well. The lemma is proved. □

5 Spaces of words

Let $\mathcal{W}$ be a set of vanishing in the group G words. An easy observation (see Lemma 5.5 below) shows that under a natural condition, spaces of words from $\mathcal{W}$ can be bounded from above by the same (up to equivalence) function as the spaces of words from a smaller set $\mathcal{W}'$. In this section, we define several sets of words (as boundary labels of several sets of diagrams) and estimate from above the spaces of words from larger sets using the upper bounds for the spaces of words from smaller sets. To apply Lemma 5.5, we will cut up a diagram Δ into two pieces such that one of the pieces belongs to a smaller set of diagrams while the perimeter of the second piece is bounded by $|\partial\Delta|$. The most difficult statement is Lemma 5.8 (separating of a one-hub subdiagram) whose proof is based on Lemma 4.9.

5.1 Spaces of boundary labels of some diagrams.

We call a disc *simple* if it has no H-cells and either all its θ-edges have labels from Θ or all of them have labels from $\hat{\Theta}$.

Lemma 5.1. *Let Δ be a diagram having exactly one hub Π. Then there is a diagram $\bar{\Delta}$ with the same boundary label as Δ such that $\bar{\Delta}$ has a simple disc subdiagram D, and the annular diagram $\Gamma = \Delta\backslash D$ is a minimal annular diagram without θ-annuli.*

Moreover, one may assume that the boundary label of D is of the form $(k_1W_1k_2W_2\dots k_LW_L)^{\pm 1}$, where $k_1W_1k_2W_2\dots k_LW_L$ is accepted by either the machine $\mathcal{S}(L)$ or by $\hat{\mathcal{S}}(L)$, and the lengths of θ-annuli in D do not exceed $N + L(space_{\mathcal{S}\cup\hat{\mathcal{S}}}(W_2))$.

Proof. We may assume that Δ is a minimal diagram. Let D_1 be a maximal disc in Δ, and denote by $\mathcal{K}_1,\dots,\mathcal{K}_L$ the maximal $k_1,\dots,k_L$-bands of D_1 starting on the hub Π. We denote by Γ_i the maximal accepted i-sector of D_1 bounded by k_i and k_{i+1} $(i = 1,\dots,L)$. By Lemmas 3.8 and 3.9(1), these sectors have no H-cells for $i \neq 1$ and each of them is a copy or a mirror copy of the 2-sector Γ_2.

Now we replace the 1-sector Γ_1 by the replica Γ_2' of Γ_2 in D_1. (To achieve this, one can make a cut along $\mathcal{K}_1$, $\mathcal{K}_2$ and the part of the boundary of Π between these k-bands, and insert two mirror copies of Γ_2' along this cut.) By the definition of replica, we obtain a modification D_2 of the disc diagram D_1 with boundary label of the form $k_1W_1k_2W_2\dots k_LW_L$, where W_1 is a mirror copy of W_2 or W_1 has no a-letters. Since $k_2W_2k_3$ is an accepted 2-sector word, the word $k_1W_1k_2W_2\dots k_LW_L$ is accepted by either the machine $\mathcal{S}(L)$ or the machine $\hat{\mathcal{S}}(L)$ by Lemma 3.1. Moreover the length of this computation does not exceed the length of the computation of $\mathcal{S}\cup\hat{\mathcal{S}}$ by the same lemma. Therefore by Lemma 3.9(2), the disc D_2 can be replaced by a disc D_3 which has no H-cells, and whose labels of θ-edges either all belong to Θ or all belong to $\hat{\Theta}$, and whose number of θ-annuli does not exceed that number for D_2.

Now, if necessary, the annular diagram $\Delta\backslash D_3$ can be replaced by a minimal diagram Γ over the group G_1. Assume that Γ has a θ-annulus $\mathcal{T}$. By Lemma 3.4, $\mathcal{T}$ surrounds the disc diagram D_3, and so D_3 can be included in a larger disc subdiagram D_4 which contains more (k_i,θ)-cells for $i \neq 1,2$ since the extensions of the $\mathcal{K}_i$-s have to cross $\mathcal{T}$. Then one can make the surgery as above and replace D_3 by a larger simple disc D_4. This procedure terminates, because we do not change the number of (k_i,θ)-cells $(i \neq 1,2)$ in the compliment of the discs when passing from D_1 to D_2 and from D_2 to D_3 and we reduce this number when passing from D_3 to D_4. (Recall that the rank of such cells is higher than the ranks of other cells in diagrams over G_1.) Thus the procedure terminates with a desired diagram Γ.

Finally, one can replace the disc by a disc corresponding to a computation of minimal space and use Lemma 3.1 to make the second claim of the lemma true.

□

Lemma 5.2. *There are positive constants c_1 and c_2 with the following property. For the boundary label $w \equiv k_1W_1\dots k_2W_l$ of the simple disc D from Lemma 5.1, there is a derivation $w \equiv w_0 \to w_1 \to \dots \to w_t = 1$ using the relations of $G(\mathcal{S}\cup\hat{\mathcal{S}},L)$ and the hub relation (3.6), such that $|w_i| \le c_1S'_{\mathcal{S}}(|W_2|) + c_2$ $(i = 0,1,\dots,t)$.*

Proof. Let us start with the words $w \equiv w(0),w(1),\dots,w(m) \equiv \Sigma_0$, written on the boundaries of the θ-annuli of D. By Lemma 5.1 and 3.1, there is $c_1 > 0$ such that $|w(i)| \le c_1S'_{\mathcal{S}}(|W_2|) + N$. Notice that $w(i)$ is written on the top of a θ annulus $\mathcal{T}$ of D and $w(i+1)$ is written on its bottom. The band $\mathcal{T}$ has

N (θ, q)-cells. For these cells the combinatorial length of their tops and bottoms differ by at most ± 1. The remaining cells are (θ, a)-cells, and their tops and bottoms have one a-edge. Therefore one can insert several elementary transformations between $w(i)$ and $w(i+1)$ corresponding to a sequential removal of the cells of $\mathcal{T}$ so that the combinatorial length of the words obtained after the refinement of our sequence does not exceed $|w(i)| + N + 2$. To complete the proof, it suffices to set $c_2 = 2N + 2$. □

Lemma 5.3. *Assume that (1) a minimal diagram Δ has no hubs and no q-bands or (2) Δ is a union of a simple disc D and a minimal annular diagram Γ over G_1 surrounding the disc subdiagram D and having no θ-annuli, and every maximal q-band of Δ starts on the hub of D. Then the sum of perimeters σ_H of all H-cells in Δ does not exceed $c_3|\partial\Delta|$ for some constant c_3 independent of Δ.*

Proof. We consider the condition (2) of the lemma only since a simplified argument works if Δ satisfies condition (1).

Two different H-cells cannot be connected by an a-band in a minimal diagram since otherwise this subdiagram could be replaced by a diagram with one H-cell and several (θ, a)-cells contrary to the minimality of the diagram. (See Lemma 3.12 (2) in [23].) A maximal a-band $\mathcal{A}$ cannot connect a-edges of the same H-cell π in a disc subdiagram by Lemma 3.12 (3) in [23], and also $\mathcal{A}$ and π cannot surround the disc D since their boundaries have no q-edges. Therefore every maximal a-band starting on an H-cell ends either on $\partial\Delta$ or on ∂D, or on a (θ, q)-cell of the annular diagram Γ. Therefore to estimate σ_H, we should give an estimate for the number $n_{\theta,q}$ of (θ, q)-cells in Γ and for the number of a-letters in the word W_1, where $k_1 W_1 \dots k_L W_L$ is the boundary label of D. (Recall that the edges of H-cells are labeled by a-letters from the alphabet $\mathcal{A}_1$ and so cannot be connected by a-bands with subpaths of ∂D labeled by $W_2, \dots, W_L$.)

Let $\mathcal{T}$ be a maximal θ-band of Γ. Since both the start and the end θ-edges of $\mathcal{T}$ must belong to $\partial\Delta$, it follows from Lemma 3.4 that $\mathcal{T}$ crosses every q-band of Γ at most once, and so has at most N (θ, q)-cells, because every maximal q-band of Γ starts on the disc D. The number of maximal θ-bands of Γ does not exceed $\partial\Delta$ since Γ has no θ-annuli. It follows that $n_{\theta,q} \le N|\partial\Delta|$.

Since we have $|W_1|_a \le |W_2|_a = |W_3|_a = \dots$ for the simple disk D, we will look for an upper estimate for $|W_2|_a$. Every maximal a-band $\mathcal{A}$ starting on the subpath p of ∂D labeled by W_2 cannot end on p by Lemma 3.4 since the word W_2 is reduced and has no θ-letters. Therefore $\mathcal{A}$ terminates on one of the two closest maximal q-bands Q and Q' starting on D or on $\partial\Delta$. The lengths of Q and Q' are at most $|\partial\Delta|$ as was explained in the previous paragraph, and so each of the sides of these q-bands has at most $|\partial\Delta|$ a-edges. Therefore $|W_2|_a \le 3|\partial\Delta|$.

Thus, $\sigma_H \le c_3|\partial\Delta|$ for the constant $c_3 = 1 + 3 + 2N$. □

Assume that a word w vanishes in the group G given by Relations (3.4), (3.5), and (3.6). Then we denote by $Space_G(w)$ the minimal number m such that there is an elementary reduction of w to the empty word such that at every step i, we have a tuple of words $(w_{i,1}, \dots, w_{i,s(i)})$ with $|w_{i,1}| + \dots + |w_{i,s(i)}| \le m$. For a set $\mathcal{W}$ of words vanishing in G (call such a set *vanishing*), we define the space function $f_{G,\mathcal{W}}(x)$ as the maximum of $Space_G(w)$ over the words $w \in \mathcal{W}$ with $|w| \le x$. (Thus we use the length $|\ |$ here unlike the length $||\ ||$ used in the Introduction.)

Now we denote by $\mathcal{W}_1$ the set of words read on the boundaries of simple discs, and we say that $w \in \mathcal{W}_2$ if w can be read on the boundary of an H-cell.

Lemma 5.4. *The function $f_{G,\mathcal{W}_2}$ is bounded from above by a function equivalent to $f_{G,\mathcal{W}_1}$.*

Proof. Assume that $|w| = n > 0$ and $w \in \mathcal{W}_2$. Since the set of generators $a_1, ..., a_m$ of H is symmetric, for every $i \le m$, we have a positive relation of the form $a_i a_{i'}$ for some $i' \le m$. By Lemma 3.2 these 2-letter relations are consequences of the relations of G. Since the set of 2-letter relations is finite, there is a constant c_4 such that one can convert w letter-for-letter to a positive word u of the same length, and the space of the corresponding derivation is $\le n + c_4$. The word u is a product of cyclic shifts of the words $\Sigma(u)$ and $\hat{\Sigma}(u)$ as was explained at the end of Subsection 3.1. Here both words $\Sigma(u)$ and $\hat{\Sigma}(u)$ belong to $\mathcal{W}_1$ and their lengths are at most $Ln + N \le (L + N)n$. Therefore for any $w \in \mathcal{W}_2$, we have $space_G(w) \le 2f_{G,\mathcal{W}_1}((L + N)n) + c_4$ which implies the statement of the lemma. □

Lemma 5.5. *Let $C > 0$. Assume that for every w from a vanishing set $\mathcal{W}$, there is a derivation $w \equiv w_0 \to w_1 \to \cdots \to w_t \equiv 1$ such that for every $i = 0, \ldots, t-1$, a cyclic shift of the word w_i is freely equal to a product of a cyclic shift of w_{i+1} and a word v_i from a vanishing set $\mathcal{W}'$, where $\max(|v_i|, |w_i|) \leq C|w|$ $(i = 0, \ldots, t-1)$. Then the function $f_{G,\mathcal{W}}$ is bounded from above by a function equivalent to $f_{G,\mathcal{W}'}$.*

Proof. The hypothesis of the lemma implies that we can apply the following series of elementary transformations to w_i. A first series of transformations replaces the word by its cyclic shift, a second series deletes/inserts mutual inverse letters operating with the words of length $\leq 2C|w|$; then we split the obtained word into a product of a cyclic shift of w_{i+1} and the word v_i; then we keep w_{i+1} unchanged and use the appropriate procedure reducing the word v_i to the empty word, and finally obtain the word w_{i+1} using cyclic shifts. Clearly, we have that the space of this procedure is at most $2C|w| + f_{G,W'}(C|w|)$. Thus, by induction on i, we have $Space_G(w) \leq 2Cn + f_{G,W'}(Cn)$ for arbitrary word $w \in \mathcal{W}$ of length at most n. The lemma is proved. □

Now we introduce the set $\mathcal{W}_3$ of boundary labels of diagrams Δ satisfying the condition of Lemma 5.3, i.e., either (1) Δ is a minimal diagram Δ having no hubs and no q-bands or (2) Δ is a union of a simple disc D and a minimal annular diagram Γ over G_1 surrounding the disc subdiagram D and having no θ-annuli, and every maximal q-band of Δ starts on the hub of D.

Lemma 5.6. *The function $f_{G,\mathcal{W}_3}$ is bounded from above by a function equivalent to $f_{G,\mathcal{W}_1 \cup \mathcal{W}_2}$.*

Proof. We will assume that a word w of length $n \geq \delta$ is the boundary label of a diagram Δ satisfying condition (2) in the definition of the set $\mathcal{W}_3$. Every cell of the annular subdiagram Γ is either an H-cell or a θ-cell. Therefore there is a sequence of diagrams $\Delta = \Delta_0, \Delta_1, \ldots, \Delta_t = D$ such that for $i = 1, \ldots, t$, the diagram Δ_i results from Δ_{i-1} after one cuts off either (a) an H-cell or (b) a rim θ-band, or an edge e such that ee^{-1} belongs to the boundary path of Δ_{i-1}. Surgery of types (b) and (c) decreases the perimeter by Lemma 4.6. Although surgery of type (a) can increase the perimeter, it follows from Lemma 5.3 that the perimeter of every diagram Δ_i is at most $(1 + c_4)n$.

Now we have a sequence $w \equiv w_0, w_1, \ldots, w_t, 1$ where $w_0, \ldots, w_t$ are the boundary labels of $\Delta_0, \Delta_1, \ldots, \Delta_t = D$, of lengths at most $(1 + c_4)n$, such that, for every $i = 0, \ldots, t$, a cyclic shift of the word w_i is a product of a cyclic shift of the word w_{i+1} and a word v_i, where v_i is either a boundary label of an H-cell of Δ or a 2-letter word aa^{-1}, or the boundary label of the simple disc D, or the boundary label of the the rim θ-band of Δ_i. In all of these cases $|v_i| \leq 2(1 + c_4)n$; in the case of the rim band, we obviously have $Space_G|v_i| \leq 2(1 + c_4)n$, and in the other cases $Space_G|v_i| \leq f_{G,\mathcal{W}_1 \cup \mathcal{W}_2}((1 + c_4)n)$. Therefore one can apply Lemma 5.5 and complete the proof. □

By definition, the set of words $\mathcal{W}_4$ contains the set $\mathcal{W}_3$ and consists of boundary labels of diagrams Δ, where either (1) Δ is a minimal diagram having no hubs or (2) Δ is a union of a simple disc D and a minimal annular diagram Γ over G_1 surrounding the disc subdiagram D and having no θ-annuli.

Lemma 5.7. *The function $f_{G,\mathcal{W}_4}$ is bounded from above by a function equivalent to $f_{G,\mathcal{W}_3}$.*

Proof. Again we assume that a word w of length $n \geq \delta$ is the boundary label of a diagram Δ satisfying condition (2) in the definition of the set $\mathcal{W}_4$. Assume that Δ has a maximal q-band $\mathcal{C}$ which does not start or terminate on the simple disc D. Then Δ is separated into 3 subdiagrams: Γ_1 contains the disc D, $\Gamma_2 = \mathcal{C}$, and Γ_3 is the remaining part of Δ. On the one hand, the lengths of the top and the bottom of $\mathcal{C}$ are equal to the number of cells m in $\mathcal{C}$ since every cell of $\mathcal{C}$ has one θ-edge and at most one a-edge on each of the sides of $\mathcal{C}$. On the other hand, each maximal θ-band of Δ crossing $\mathcal{C}$ must start and terminate on $\partial\Delta$. This implies that the perimeters of each of the subdiagrams Γ_1, Γ_2, and Γ_3 are at most $2n$. (Here we use the definition of length and take into account that the band $\mathcal{C}$ starts and ends on q-edges.)

Therefore there is a sequence of diagrams $\Delta = \Delta_0, \Delta_1, \ldots, \Delta_t$ of perimeters $\leq 2n$ such that for $i = 1, \ldots, t-1$, the diagram Δ_i results from Δ_{i-1} after one cut off either (a) a subdiagram without q-bands or (b) a q-band, and Δ_t has no maximal q-bands except for those starting/terminating on the hub of D. Let $w \equiv w_0, w_1, \ldots, w_t$ be the boundary labels of these diagrams. Every word w_i $(i = 0, \ldots, t)$ of the series $w \equiv w_0, w_1, \ldots, w_t, w_{t+1} = 1$ (or its cyclic shift) is a product of a cyclic shift of the word w_{i+1} and a word v_i, where v_i either belongs to $\mathcal{W}_3$ or it is the boundary label of a q-band. In all of these

cases $|v_i| \le 2n$, and in the later case we obviously have $Space_G|v_i| \le 2n$. In the former case we have $space_G|v_i| \le f_{G,\mathcal{W}_3}(2n)$. To complete the proof, we apply Lemma 5.5. □

Lemma 5.8. *Let $\mathcal{W}_5$ be the set of all words vanishing in G. The function $f_{G,\mathcal{W}_5}(n)$ is bounded from above by a function equivalent to $f_{G,\mathcal{W}_3\cup\mathcal{W}_4}(n)$.*

Proof. Let $w = 1$ in G and $|w| = n > 0$. If $w = 1$ in G_1, then $Space_G(w) \le Space_{G,\mathcal{W}_3}(w)$ by Lemma 5.7. Otherwise the minimal diagram Δ with boundary label w has $t \ge 1$ hubs, and by Lemma 4.9, a cyclic shift of the word $w \equiv w_0$ is a product of a word v_1 written on the boundary of a diagram Γ having one hub, and a word w_1 which is a boundary label of a diagram with $t-1$ hubs, and $|w_1| \le n, |v_1| \le 2n$. By Lemma 5.1, $v_1 \in \mathcal{W}_4$. Now induction on t gives a series of words $w_0, w_1, \dots, w_t = 1$ and words $v_1, \dots, v_t$ satisfying the conditions of Lemma 5.5 with $C = 2$, and our statement follows from that lemma. □

5.2 Proofs of main statements.

Proof of Theorem 1.2. Suppose H is a group with a finite generating set $\{a_1, \dots, a_m\}$ and the word problem for H is solvable by a DTM with space complexity $f(n)$. As we mentioned in the beginning of Subsection 3.1, one can double the set of generators so that every a_i^{-1} becomes equal in H to some a_j. Hence the set of positive words vanishing in H is a set of defining relations for H, and it is recognizable by a DTM M with space complexity $f(n)$. By Lemma 2.10 (a), there is an S-machine $\mathcal{S}$ recognizing the same language, and the generalized space complexity $S'_{\mathcal{S}}(n)$ of $\mathcal{S}$ is equivalent to $f(n)$. The group G constructed on the basis of $\mathcal{S}$ in Subsection 3.1 is finitely presented and contains H as a subgroup by Corollary 3.1. The consecutive application of Lemmas 5.8, 5.7, 5.6, 5.4, and 5.2 results in the inequality $f_{G,\mathcal{W}_5}(n) \preceq S'_{\mathcal{S}}(n)$. Note that the space functions $f_{G,\mathcal{W}_5}(n)$ and $s_G(n)$ of G are equivalent since the lengths functions $||*||$ and $|*|$ satisfy inequalities $\delta||w|| \le |w| \le ||w||$ for every word w. Hence $s_G(n) \preceq S'_{\mathcal{S}}(n)$. Since $S'_{\mathcal{S}}(n) \sim f(n)$ by Lemma 2.10 (2), we have $s_G(n) \preceq f(n)$.

To invert this inequality and to obtain the second claim of Theorem 1.2, we first note that $s_G(n) \succeq \log d_G(n)$, where $d_G(n)$ is the Dehn function for G. (Indeed, up to equivalence, the length t of a rewriting $W_0 = (w_0) \to \dots \to W_t = (1)$ without repetitions does not exceed $\exp(\max_{i=0}^{t} ||W_i||)$; see also Theorem C in [7]). Therefore it suffices to show that $f(n) \preceq \log d_G(n)$. By Lemma 2.4(3), $S'_{M'}(n) \sim f(n)$, and by Lemma 2.9 (4), $T'_{\mathcal{S}}(n) \succeq \exp(S'_{M'}(n))$. Thus it remains to explain that $d_G(n) \succeq T'_{\mathcal{S}}(n)$.

Let W be a word accepted by $\mathcal{S}$, such that $1 \le |W|_a \le n$ and $time_{\mathcal{S}}(W) = T'_{\mathcal{S}}(n)$. Then the word $V \equiv k_1 W_1 \dots k_L W_L$ (where W_i-s are copies or mirror copies of W) is accepted by $\mathcal{S}(L)$ and therefore it is conjugate to the word Σ_0 (see subsection 3.1). Hence $V = 1$ in G. Let Δ be a minimal diagram over G with boundary label V. Since $\partial\Delta$ has only one k_1-edge, the maximal k_1-band starting on this k_1-edge must end on the boundary of a hub. Hence Δ has a hub Π satisfying the condition of Lemma 3.11. If one removes Π together with the bands $\mathcal{B}_1, \dots, \mathcal{B}_{L-3}$ and subdiagrams Γ_i $(i = 1, \dots, L-4)$, then the remaining diagram Δ' has at most $L - (L-3) + 3 = 6$ k-edges. It follows from Lemma 3.11 that Δ' has no hubs since $6 < L - 3$. Thus Δ has exactly one hub.

Obviously, every k_i-band starting on $\partial\Pi$ ends on $\partial\Delta$ $(i = 1, \dots, L)$, and so we can consider the 2-sector Γ of Δ bounded by k_2- and k_3-bands. By Lemma 3.9(1), the number of maximal θ-bands of Γ is equal to the length of a computation of $\mathcal{S}(L) \cup \hat{\mathcal{S}}(L)$ accepting the word W. By Lemma 3.1, we can replace $\mathcal{S}(L) \cup \hat{\mathcal{S}}(L)$ by $\mathcal{S}$ in the previous phrase, and so $area(\Delta) > T'_{\mathcal{S}}(n)$. Since $||V|| \le C|W|_a$ for a constant C, we have $d_G(Cn) > T'_{\mathcal{S}}(n)$, and the required lower bound is obtained. □

Proof of Theorem 1.4. Let M be a DTM with space complexity $f(n)$. Then, as in the proof of Theorem 1.2, we can construct an S-machine $\mathcal{S}$ with $S'_{\mathcal{S}}(n) \sim f(n)$ and $T'_{\mathcal{S}}(n) \sim \exp f(n)$. But now we simplify the construction of the group G: It is defined by the relations associated with the machine $\mathcal{S}(L)$ only and the hub relation (there is no $\hat{\mathcal{S}}(L)$ now, and so we have no H-relations at all). The (simplified) proof of Theorem 1.2 works in this setting and therefore $s_G(n) \sim f(n)$ and $\log d_G(n) \sim f(n)$. Finally, by Lemma 3.9 (1,2) and by the argument exploited in the proof of Theorem 1.2, an input word u of M is accepted iff the word $\Sigma(u)$ vanishes in G; this completes the proof. □

Proof of Corollary 1.6. Let an NTM have FSC space complexity $f(n)$ and solve the word problem in a finitely generated group H. Then by Savitch's Theorem (see [9], Theorem 1.30), there is a DTM which solves the same problem with space $\sim f(n)^2$. It remains to refer to Theorem 1.2. □

Remark 5.9. Corollary 1.6 has a problem-reduction version. If a DTM M has an FSC space complexity function $f(n)$, then one can construct a finitely presented group $G = G(M)$ according to Theorem 1.4, and so G has a space function equivalent to $f(n)$. By Proposition 1.1, the word problem of G is solvable by an NTM of space complexity $f(n)$, and so it is solvable by a DTM with space $f(n)^2$. Therefore Corollary 1.5 can be extended to say that many space complexity classes have space complete word problem for some finitely presented group (depending on the class) with space function equivalent to the given space complexity $f(n)$.

We need one more lemma to prove Corollary 1.7. This is a version of Savitch's theorem (see Theorem 1.30 in [9]), but instead of simulation of the work of an NTM by a DTM, now we need a DTM computing the space complexity of a given NTM.

Lemma 5.10. *Let M be an NTM with space complexity $S(n)$ bounded from above by an FSC function $f(n)$. Then there is a DTM M_0 such that (1) M_0 computes $S(n)$, i.e., for any input $n \in \mathbb{N}$ given in binary, it computes $S(n)$; (2) the space complexity $S_{M_0}(n)$ is $O(f(n)^2)$.*

Proof. Without loss of generality, we may assume that M satisfies the $\vec{s}_{10}$ condition.

If u is an accepted input word for M and $||u|| \leq n$, then the time of any computation (without repetitions) of space $\leq f(n)$ accepting u is at most $\leq 2^{cf(n)}$, for some integer $c > 0$ since all configurations in this computation are of length $\leq f(n) + c_0$ for a constant c_0. So the goal for the DTM M' we want to define, is to find a computation C of M of minimal space and of length at most $2^{cf(n)}$ which connects the input configuration and the accept configuration of M, and then to compute the space of this computation. (If such a computation exists; otherwise M' says that $u \notin \mathcal{L}_M$.) Indeed, the required machine M_0 will examine all words u with $||u|| \leq n$ in lexicographical order, it will switch on M' for every such input word u, and it will compare the spaces $space_M(u)$ of the u-s on an additional tape keeping only the maximal one after every return.

For arbitrary words w and w' and $k \in \mathbb{N}$, define predicates $reach_n(w, w', k)$ to mean that w and w' are configurations of M and there is a computation $w \to \cdots \to w'$ with time $\leq k$ and with space $\leq f(n) + c_0$. By this definition, M accepts an input word u of combinatorial length $\leq n$ iff $reach_n(w_0, w_f, 2^{cf(n)})$, where $w_0 \equiv w(u)$ is the unique input configuration on input u and w_f is the unique accept configuration of M.

Note that $reach_n(w, w', k + j)$ iff $(\exists w'')(reach_n(w, w'', k)$ *and* $reach_n(w'', w', j))$ and the minimal space of computations $w \to \cdots \to w'$ is the minimum over all w'' of the maximums of minimal spaces for $w \to \cdots \to w''$ and for $w'' \to \cdots \to w'$. This observation leads to the following (slight) modification of Savitch's machine.

- Given $n \in \mathbb{N}$, compute $2^{cf(n)}$ (in binary) using that $f(n)$ is an FSC function.
- Then, $reach_n(w, w', 1)$ is true if $w \equiv w'$ or w transforms into w' under the application of a single command of M. The space of the computation $w \to w'$ is $\max(|w|_a, |w'|_a)$.
- If $k \geq 2$, then for all possible configurations w'' of M with length $\leq f(n) + c_0$, compute whether it is true that $reach_n(w, w'', [(k+1)/2])$ and $reach_n(w'', w', [(k+1)/2)]$. Set $reach_n(w, w', k)$ to be true iff such w'' exists. Find the minimal space of computations $w \to \cdots \to w'$ of length $\leq k$ using the information on the minimal space of computations $w \to \cdots \to w''$ and $w'' \to \cdots \to w'$ of length $\leq [(k+1)/2]$.

It is easy to see that the constructed machine M' computes, in particular, the space of any accepted input word u of length $\leq n$. Passing from k to $[(k+1)/2]$, we need additional space to store the information on k, w, w', on the current w'', and afterwards, on the minimal space of computations $w \to \cdots \to w'$ of length $\leq k$. Clearly, this additional space is $O(f(n))$, and since we start with $k = 2^{cf(n)}$, we divide k by 2 $cf(n)$ times, so the total space used by M' and by M_0 is $O(f(n)^2)$.

□

Proof of Corollary 1.7. Assume that α is computable with space $\leq 2^{2^m}$. It follows that for $m = [\log_2 \log_2 n]$ we can recursively compute binary rationals α_m such that

$$|\alpha - \alpha_m| = O(2^{-m}) = O((\log_2 n)^{-1}) \tag{5.8}$$

and the space of the computation of α_m is at most n. In addition, one may assume that the number of digits in the binary expansion of α_m is $O(m)$. Therefore the computation of $[\log_2 n]$ (in binary) and of the product $\alpha_m[\log_2 n]$ needs space at most $O((\log_2 n)^2)$. Then we rewrite the binary presentation of $[\alpha_m[\log_2 n]]$ in unary (as a sequence of 1-s). This well-known rewriting (e.g., see p.352 in [27]) has space complexity of the form $[\alpha_m[\log_2 n]] + O(1)$. One more rewriting of this type applied to the unary presentation of $[\alpha_m[\log_2 n]]$, will have space complexity of the form $2^{[\alpha_m[\log_2 n]]} + O(1)$. Using (5.8), we can present this function as

$$2^{\alpha[\log_2 n]+O(1)} + O(1) \sim 2^{\alpha[\log_2 n]} \sim [n^\alpha]$$

Thus the subsequent application of the above mentioned DTM-s has space complexity equivalent to n^α, and we can apply Corollary 1.4 to obtain a finitely presented group with space function equivalent to n^α.

Now assume that a function $[n^\alpha]$ is equivalent to a space function of a finitely presented group G. Then by Proposition 1.1, there is an NTM M whose space complexity $S_M(n)$ is equivalent to $[n^\alpha]$, that is

$$c_1 n^\alpha < S_M(n) < c_2 n^\alpha \tag{5.9}$$

for some positive c_1, positive integer c_2, and every sufficiently large n. In particular, we have $S_M(n) < c_2 n^d$ for some integer d and every n. Since $c_2 n^d$ is an FSC function, we may apply Lemma 5.10, and obtain a DTM M_0 computing the function $S_M(n)$ with space $O(n^{2d})$. Hence M_0 computes the function $S_M(2^{2^m})$ of m with space $O((2^{2^m})^{2d})$. This space is less than $2^{2^{m+c_3}}$ for some c_3. Hence for some $c_4 \in \mathbb{N}$, M_0 computes the function $S_M(2^{2^{m-c_4}})$ with space at most $2^{2^{m-1}}$.

Let us plug $n = 2^{2^{m-c_4}}$ into Inequalities (5.9) and then take $\log_2$ of the terms. We obtain

$$\lambda_1 + \alpha 2^{m-c_4} \leq \log_2 S_M(2^{2^{m-c_4}}) \leq \lambda_2 + \alpha 2^{m-c_4}$$

where $\lambda_i = \log_2 c_i$ $(i = 1, 2)$. It follows that

$$|\alpha - 2^{-m+c_4} \log_2 S_M(2^{2^{m-c_4}})| < c 2^{-m+c_4} = O(2^{-m}) \tag{5.10}$$

where $c = \max(|\lambda_1|, |\lambda_2|)$. Recall that $S_M(2^{2^{m-c_4}}) \leq c_2 (2^{2^{m-c_4}})^d \leq 2^{2^{m-1}}$ and so this number has at most $2^{m-1} + 1$ binary digits. Therefore the real numbers $\log_2(S_M(2^{2^{m-c_4}}) 2^{-m+c_4})$ are computable with error $O(2^{-m})$ and space $O(2^{2^{m-1}})$. Now it follows from (5.10) that the real number α is computable with space 2^{2^m}. □

Acknowledgment.

The author is grateful to J.-C. Birget, M. Bridson, T.Davis, M. Elder, S.V.Ivanov, T.Riley, M. Sapir, and the anonymous referee for useful discussions, comments, and criticism.

References

[1] G. Baumslag, A non-cyclic one-relator group all of whose finite quotients are cyclic. J. Austral. Math. Soc., 10 (1969), 497-498.

[2] J.-C. Birget, Time-complexity of the word problem for semigroups and the Higman embedding theorem, Internat. J. Algebra Comput. 8 (1998), 235-294.

[3] J.-C. Birget, Functions on groups and computational complexity, Internat. J. Algebra Comput., 14 (2004), no. 4, 409–429.

[4] J.-C. Birget, J.-C., A. Yu. Olshanskii, E. Rips, M. V. Sapir, Isoperimetric functions of groups and computational complexity of the word problem. Ann. of Math. (2) 156 (2002), no. 2, 467–518.

[5] N.Brady and M.Bridson, There is only one gap in the isoperimetric spectrum, Geometric and Functional Analysis, 10 (2000), 1053-1070.

[6] N. Brady, T.Riley, and H.Short, The geometry of the word problem for finitely generated groups, Advanced Courses in Mathematics, CRM Barselona, Birkhauser-Verlag, Basel, 2007, x+206 p.p.

[7] M.R. Bridson, T.R. Riley, Free and fragmenting filling length, Journal of Algebra, 307(1) (2007), 171-190.

[8] D.E. Cohen, K.Madlener, and F.Otto, Separating the intrinsic complexity and the derivational complexity of the word problem for finitely presented groups, Math. Logic Quart, 39, no. 2 (1993), 143-157.

[9] Ding-Zhu Du, Ker-I Ko, Theory of Computational Complexity, Wiley-Interscience Publ., N.Y.,2000, 512 p.p.

[10] S. M. Gersten, Dehn functions and l1-norms of finite presentations. Algorithms and Classification in Combinatorial Group Theory, Springer, Berlin, 1992, 195225.

[11] S.M.Gersten, Isoperimetric and isodiametric functions. In G.Niblo and M.Roller editors, Geometric group theory I, Lecture Notes of LMS, 181, Camb. Univ. Press, 1993.

[12] S.M.Gersten, T.R.Riley, Filling length in finitely presentable groups, Geometricae Dedicata, 92(1) (2002), 41-58.

[13] M.Gromov, Hyperbolic groups, in: Essays in Group Theory (S.M.Gersten, ed.), M.S.R.I. Pub. 8, Springer, 1987, 75–263.

[14] M.Gromov, Asymptotic invariants of infinite groups, in: Geometric Group Theory. Vol. 2 (G.A.Niblo and M.A.Roller, eds.), London Math. Soc. Lecture Notes Ser., 182 (1993), 1–295.

[15] V.Guba, M.Sapir, On Dehn functions of free products of groups, Proc. Amer. Math. Soc. 127 (1999), 1885–1891.

[16] R.Lipton, Y.Zalcstein, Word problems solvable in logspace, J. Assoc. Comput. Mach., 24 (1977), no. 3, 522–526.

[17] R.C. Lyndon, P.E. Schupp, Combinatorial Group Theory, Springer–Verlag, 1977.

[18] K.Madlener, F.Otto, Pseudo-natural algorithms for the word problem for finitely presented monoids and groups, J. Symbolic Computation 1(1985), 383-418.

[19] A. G. Miasnikov, A. Ushakov, and Dong Wook Won, Word problem in Baumslag-Gersten group is polynomial time decidable, to appear.

[20] A.Yu.Olshanskii, On the subgroup distortion in finitely presented groups, Matem. Sbornik, 188 (1997), N 11, 73-120 (in Russian).

[21] A.Yu.Olshanskii and M.V.Sapir, Length and area functions in groups and quasi-isometric Higman embeddings, Intern. J. Algebra and Comput., 11 (2001), no. 2 , 137-170.

[22] A.Yu.Olshanskii and M.V.Sapir, Non-amenable finitely presented torsion-by-cyclic groups, Publ. Math. IHES, 96 (2003), no. 6, p.p. 43 - 169

[23] A.Yu.Olshanskii and M.V.Sapir, Conjugacy problem and Higman embeddings, "Memoirs of the AMS" 170(2004), no. 804 p.p. vii+131

[24] A.Yu.Olshanskii and M.V. Sapir, Groups with small Dehn functions and bipartite chord diagrams, Geometric and Functional Analysis, 16 (2006), 1324-1376

[25] A.N.Platonov, Isoperimetric function of the Baumslag-Gersten group, (Russian) Vestnik Moskov. Univ. Ser. I Mat. Mekh. (2004), pp. 1217.

[26] J.Rotman, An introduction to the theory of groups, 3d edition, Allyn and Bacon Inc., Boston, Mass, 1984.

[27] M. V. Sapir, J. C. Birget, E. Rips, Isoperimetric and isodiametric functions of groups, Annals of Mathematics, 157, 2(2002), 345-466.

[28] B.A.Trakhtenbrot, On the complexity of reduction algorithms in Novikov - Boone constructions, Algebra i Logika 8(1969), no. 1, pp. 93-128; English translation in: Algebra and Logic, 8(1969), no. 1, pp. 50-71.

[29] M.K.Valiev, On the complexity of the identity problem for finitely defined groups, Algebra i Logika 8(1969), no. 1, pp. 5-43; English translation in: Algebra and Logic, 8(1969), no. 1, pp. 2-21.